

Spectrum of a non-self-adjoint operator associated with the periodic heat equation

Marina Chugunova, Dmitry Pelinovsky*

Department of Mathematics, McMaster University, Hamilton, Ontario, Canada, L8S 4K1

Received 26 February 2007

Available online 23 December 2007

Submitted by M. Nakao

Abstract

We study the spectrum of the linear operator $L = -\partial_\theta - \epsilon \partial_\theta (\sin \theta \partial_\theta)$ subject to the periodic boundary conditions on $\theta \in [-\pi, \pi]$. We prove that the operator is closed in $L^2_{\text{per}}([-\pi, \pi])$ with the domain in $H^1_{\text{per}}([-\pi, \pi])$ for $|\epsilon| < 2$, its spectrum consists of an infinite sequence of isolated eigenvalues and the set of corresponding eigenfunctions is complete. By using numerical approximations of eigenvalues and eigenfunctions, we show that all eigenvalues are simple, located on the imaginary axis and the angle between two subsequent eigenfunctions tends to zero for larger eigenvalues. As a result, the complete set of linearly independent eigenfunctions does not form a basis in $L^2_{\text{per}}([-\pi, \pi])$.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Spectrum of a non-self-adjoint operator; Advection–diffusion equation with periodic coefficients; Completeness and basis of eigenfunctions; Numerical approximation of eigenvalues and eigenfunctions

1. Introduction

We address the Cauchy problem for the periodic heat equation

$$\begin{cases} \dot{h} = -h_\theta - \epsilon (\sin \theta h_\theta)_\theta, & t > 0, \\ h(0) = h_0, \end{cases} \quad (1.1)$$

subject to the periodic boundary conditions on $\theta \in [-\pi, \pi]$. This model was derived in the context of the dynamics of a thin viscous fluid film on the inside surface of a cylinder rotating around its axis in [3]. Extension of the model to the three-dimensional motion of the film was reported in [4].

The parameter ϵ is small for applications in fluid dynamics [3] and our main results correspond to the interval $|\epsilon| < 2$ in accordance to these applications. For any $\epsilon > 0$, the Cauchy problem for the heat equation (1.1) on the half-interval $\theta \in [0, \pi]$ is generally ill-posed [13] and it is naturally to expect that the Cauchy problem remains ill-posed on the entire interval $\theta \in [-\pi, \pi]$. The authors of the pioneer work [3] used a heuristic asymptotic solution to suggest that the growth of “explosive instabilities” might occur in the time evolution of the Cauchy problem (1.1).

* Corresponding author.

E-mail address: dmpeli@math.mcmaster.ca (D. Pelinovsky).

Nevertheless, in a contradiction with the picture of explosive instabilities, only purely imaginary eigenvalues were discovered in the discrete spectrum of the associated linear operator

$$L = -\epsilon \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) - \frac{\partial}{\partial \theta}, \tag{1.2}$$

acting on sufficiently smooth periodic functions $f(\theta)$ on $\theta \in [-\pi, \pi]$. Various approximations of eigenvalues were obtained in [3] by two asymptotic methods (expansions in powers of ϵ and the WKB method) and by three numerical methods (the Fourier series approximations, the pseudospectral method, and the Newton–Raphson iterations). The results of the pseudospectral method were checked independently in [17] (see pp. 124–125 and 406–408). It is seen both in [3] and [17] that the level sets of the resolvent $(\lambda - L)^{-1}$ form divergent curves to the left and right half-planes and, while true eigenvalues lie on the imaginary axis, eigenvalues of the truncated Fourier series may occur in the left and right half-planes of the spectral plane. This distinctive feature was interpreted in [3] towards the picture of growth of disturbances and the phenomenon of explosive instability.

One more question raised in [3] was about the validity of the series of eigenfunctions associated to the purely imaginary eigenvalues of the operator L for $\epsilon \neq 0$. Although various initial conditions h_0 were decomposed into a finite sum of eigenfunctions and the error decreased with a larger number of terms in the finite sum, the authors of [3] conjectured that the convergence of the series depended on the time variable and “even though the series converges at $t = 0$, it may diverge later.” This conjecture would imply that the eigenfunctions of L for $\epsilon \neq 0$ do not form a basis of functions in the space $L^2_{\text{per}}([-\pi, \pi])$ unlike the harmonics of the complex Fourier series associated with the operator L for $\epsilon = 0$.

In this paper, we prove that the operator L is closed in $L^2_{\text{per}}([-\pi, \pi])$ with a domain in $H^1_{\text{per}}([-\pi, \pi])$ for $|\epsilon| < 2$, such that the spectrum of the eigenvalue problem

$$-\epsilon \frac{d}{d\theta} \left(\sin \theta \frac{df}{d\theta} \right) - \frac{df}{d\theta} = \lambda f, \quad f \in H^1_{\text{per}}([-\pi, \pi]), \tag{1.3}$$

is well-defined. Here and henceforth, we denote

$$H^1_{\text{per}}([-\pi, \pi]) = \{f \in H^1([-\pi, \pi]): f(\pi) = f(-\pi)\}. \tag{1.4}$$

Furthermore, we prove that the residual and continuous spectra of the spectral problem (1.3) are empty and the eigenvalues of the discrete spectrum accumulate at infinity along the imaginary axis. We further prove completeness of the series of eigenfunctions associated to all eigenvalues of the purely discrete spectrum of L in $L^2_{\text{per}}([-\pi, \pi])$. Using the numerical approximations of eigenvalues and eigenfunctions of the spectral problem (1.3), we show that all eigenvalues of L are simple, located at the imaginary axis, and the angle between two subsequent eigenfunctions tends to zero for larger eigenvalues. As a result, the complete set of linearly independent eigenfunctions does not form a basis in $L^2_{\text{per}}([-\pi, \pi])$ and hence it cannot be used to solve the Cauchy problem associated with the heat equation (1.1).

The paper is structured as follows. Sections 2–5 present main results of our studies. Properties of the operator L are analyzed in Section 2. Eigenvalues of the operator L are characterized in Section 3. Section 4 presents numerical approximations of eigenvalues and eigenfunctions of the spectral problem (1.3). Section 5 discusses the Cauchy problem for the heat equation (1.1). Appendices A and B give supplementary material to the main text. An extension of the spectral problem (1.3) into a self-adjoint problem in a weighted L^2 -space is reported in Appendix A. Eigenvalues of the operator L are shown to be resonance poles of a linear Schrödinger operator on an infinite line in Appendix B.

2. General properties of the linear operator L

It is obvious that the operator L is densely defined in $L^2_{\text{per}}([-\pi, \pi])$ on the space of smooth functions with periodic boundary conditions. However, the operator L is not closed in $L^2_{\text{per}}([-\pi, \pi])$ if the functions are infinitely smooth. We therefore prove in Lemma 1 that the operator L admits a closure in $L^2_{\text{per}}([-\pi, \pi])$ with a domain in $H^1_{\text{per}}([-\pi, \pi])$. Eigenfunctions and eigenvalues of the spectral problem (1.3) are studied in Lemmas 2 and 3. The absence of the residual and continuous spectra of operator L is proved in Lemmas 4 and 5.

Lemma 1. *The operator L admits a closure in $L^2_{\text{per}}([-\pi, \pi])$ for $|\epsilon| < 2$ with $\text{Dom}(L) \subset H^1_{\text{per}}([-\pi, \pi])$.*

Proof. According to Lemma 1.1.2 in [5], if an operator has a non-empty spectrum in a proper subset of a complex plane, then it must be closed. The operator L has a non-empty spectrum in $L^2_{\text{per}}([-\pi, \pi])$ since $\lambda = 0$ is an eigenvalue with the eigenfunction $f_0(\theta) = 1 \in L^2_{\text{per}}([-\pi, \pi])$. We should show that there exists at least one regular point $\lambda_0 \in \mathbb{C}$, such that

$$\forall f \in H^1_{\text{per}}([-\pi, \pi]): \quad \|(L - \lambda_0 I)f\|_{L^2_{\text{per}}([-\pi, \pi])} \geq k_0 \|f\|_{L^2_{\text{per}}([-\pi, \pi])} \tag{2.1}$$

for some $k_0 > 0$. By using straightforward computations, we obtain

$$(f', Lf) = - \int_{-\pi}^{\pi} (1 + \epsilon \cos \theta) |f'|^2 d\theta - \epsilon \int_{-\pi}^{\pi} \sin \theta \tilde{f}' f'' d\theta, \tag{2.2}$$

where $(g, f) = \int_{-\pi}^{\pi} \bar{g}(\theta) f(\theta) d\theta$ is a standard inner product in L^2 . If $f \in H^1_{\text{per}}([-\pi, \pi])$, then

$$\text{Re}(f', f) = 0, \quad \text{Re}(f', Lf) = - \int_{-\pi}^{\pi} \left(1 + \frac{\epsilon}{2} \cos \theta\right) |f'|^2 d\theta, \tag{2.3}$$

such that for any $\lambda_0 \in \mathbb{R}$ it is true that

$$|\text{Re}(f', (L - \lambda_0 I)f)| \geq \left(1 - \frac{|\epsilon|}{2}\right) \|f'\|_{L^2_{\text{per}}([-\pi, \pi])}^2.$$

Any periodic function $f \in H^1_{\text{per}}([-\pi, \pi])$ is represented by $f_0 + \tilde{f}(\theta)$, where $f_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\theta) d\theta$ and $\tilde{f}(\theta)$ belongs to the space

$$H_0 = \left\{ \tilde{f} \in H^1_{\text{per}}([-\pi, \pi]): \int_{-\pi}^{\pi} \tilde{f}(\theta) d\theta = 0 \right\}. \tag{2.4}$$

By using the Cauchy–Schwarz inequality, we obtain

$$|\text{Re}(\tilde{f}', (L - \lambda_0 I)\tilde{f})| \leq |(\tilde{f}', (L - \lambda_0 I)\tilde{f})| \leq \|\tilde{f}'\|_{L^2_{\text{per}}([-\pi, \pi])} \|(L - \lambda_0 I)\tilde{f}\|_{L^2_{\text{per}}([-\pi, \pi])},$$

such that, for any $|\epsilon| < 2$,

$$\|(L - \lambda_0 I)\tilde{f}\|_{L^2_{\text{per}}([-\pi, \pi])}^2 \geq \left(1 - \frac{|\epsilon|}{2}\right)^2 \|f'\|_{L^2_{\text{per}}([-\pi, \pi])}^2.$$

According to the Neumann–Poincare inequality on $[-\pi, \pi]$, we have

$$\|f\|_{L^2_{\text{per}}([-\pi, \pi])}^2 \leq 4\pi^2 \|f'\|_{L^2_{\text{per}}([-\pi, \pi])}^2 + \frac{1}{2\pi} \left(\int_{-\pi}^{\pi} f(\theta) d\theta \right)^2, \tag{2.5}$$

such that

$$\begin{aligned} \|(L - \lambda_0 I)f\|_{L^2_{\text{per}}([-\pi, \pi])}^2 &= 2\pi \lambda_0^2 f_0^2 + \|(L - \lambda_0 I)\tilde{f}\|_{L^2_{\text{per}}([-\pi, \pi])}^2 \\ &\geq 2\pi \lambda_0^2 f_0^2 + \left(1 - \frac{|\epsilon|}{2}\right)^2 \|f'\|_{L^2_{\text{per}}([-\pi, \pi])}^2 \geq \lambda_0^2 \|f\|_{L^2_{\text{per}}([-\pi, \pi])}^2, \end{aligned}$$

if $\lambda_0 = \frac{1}{2\pi} (1 - \frac{|\epsilon|}{2})$. Thus, the estimate (2.1) holds for $|\epsilon| < 2$ with $k_0 = \lambda_0$. \square

Remark 1. The formal adjoint of L in $L^2_{\text{per}}([-\pi, \pi])$ is $L^* = -\epsilon \partial_{\theta} (\sin \theta \partial_{\theta}) + \partial_{\theta}$. According to Lemma 1.2.1 in [5], the operator L^* , similarly to operator L , also admits a closure in $L^2_{\text{per}}([-\pi, \pi])$ with $\text{Dom}(L^*) \subset H^1_{\text{per}}([-\pi, \pi])$ for $|\epsilon| < 2$.

Lemma 2. Let λ be an eigenvalue of the spectral problem $Lf = \lambda f$ with an eigenfunction $f \in H^1_{\text{per}}([-\pi, \pi])$. Then,

- (i) $-\lambda, \bar{\lambda}$ and $-\bar{\lambda}$ are also eigenvalues of the spectral problem $Lf = \lambda f$ with the eigenfunctions $f(-\theta), \bar{f}(\theta)$ and $\bar{f}(-\theta)$ in $H^1_{\text{per}}([-\pi, \pi])$.
- (ii) λ is also an eigenvalue of the adjoint spectral problem $L^* f^* = \lambda f^*$ with the eigenfunction $f^* = f(\pi - \theta)$ in $H^1_{\text{per}}([-\pi, \pi])$.
- (iii) λ is a simple isolated eigenvalue of $Lf = \lambda f$ if and only if $(f^*, f) \neq 0$.

Proof. (i) Due to inversion $\theta \rightarrow -\theta$, the spectral problem (1.3) transforms to itself with the transformation $\lambda \rightarrow -\lambda$. Due to the complex conjugation, it transforms to itself with $\lambda \rightarrow \bar{\lambda}$.

(ii) Due to the transformation $\theta \rightarrow \pi - \theta$, the spectral problem (1.3) transforms to the adjoint problem $L^* f = \lambda f$ with the same eigenvalue.

(iii) The assertion follows by the Fredholm Alternative Theorem for isolated eigenvalues. \square

Lemma 3. Let λ be an eigenvalue of the spectral problem (1.3) with the eigenfunction $f \in H^1_{\text{per}}([-\pi, \pi])$. Then,

$$\text{Re}(\lambda) = \epsilon \frac{(f', \sin \theta f')}{(f, f)}, \quad i \text{Im}(\lambda) = \frac{(f', f)}{(f, f)}, \tag{2.6}$$

and $\text{Im}(\lambda) \neq 0$ except for a simple zero eigenvalue $\lambda = 0$.

Proof. By constructing the quadratic form for $f \in H^1_{\text{per}}([-\pi, \pi])$, we obtain

$$(f, Lf) = \epsilon \int_{-\pi}^{\pi} \sin \theta |f'|^2 d\theta - \int_{-\pi}^{\pi} \bar{f} f' d\theta, \tag{2.7}$$

where the second term is purely imaginary since

$$f \in H^1_{\text{per}}([-\pi, \pi]): \int_{-\pi}^{\pi} \bar{f} f' d\theta = |f(\theta)|^2 \Big|_{\theta=-\pi}^{\theta=\pi} - \int_{-\pi}^{\pi} \bar{f} f' d\theta = - \int_{-\pi}^{\pi} \overline{\bar{f} f'} d\theta. \tag{2.8}$$

Moreover, the equality (2.3) can be rewritten in the form

$$i \text{Im}(\lambda)(f', f) = \text{Re}(f', Lf) = - \int_{-\pi}^{\pi} \left(1 + \frac{\epsilon}{2} \cos \theta\right) |f'(\theta)|^2 d\theta \leq - \left(1 - \frac{|\epsilon|}{2}\right) \|f'\|_{L^2}^2, \tag{2.9}$$

where the right-hand side is negative if $|\epsilon| < 2$ and $f(\theta)$ is not constant on $\theta \in [-\pi, \pi]$. Therefore, $(f', f) \neq 0$ and $\text{Im}(\lambda) \neq 0$. Finally, the constant eigenfunction $f(\theta) = 1$ corresponds to the eigenvalue $\lambda = 0$ and it is a simple eigenvalue since $(f^*, f) \neq 0$, where $f^*(\theta) = f(\pi - \theta) = 1$ is an eigenfunction of the adjoint operator L^* for the same eigenvalue $\lambda = 0$. \square

Lemma 4. The residual spectrum of the operator L is empty.

Proof. By a contradiction, assume that λ belongs to the residual part of the spectrum of L such that $\text{Ker}(L - \lambda I) = \emptyset$ but $\text{Range}(L - \lambda I)$ is not dense in $L^2_{\text{per}}([-\pi, \pi])$. Let $g \in L^2_{\text{per}}([-\pi, \pi])$ be orthogonal to $\text{Range}(L - \lambda I)$, such that

$$\forall f \in L^2([-\pi, \pi]): 0 = (g, (L - \lambda I)f) = ((L^* - \bar{\lambda} I)g, f).$$

Therefore, $(L^* - \bar{\lambda} I)g = 0$, that is $\bar{\lambda}$ is an eigenvalue of L^* . By Lemma 2(ii), $\bar{\lambda}$ is an eigenvalue of L and by Lemma 2(i), λ is also an eigenvalue of L . Hence λ cannot be in the residual part of the spectrum of L . \square

Lemma 5. The continuous spectrum of the operator L is empty.

Proof. According to Theorem 4 in [8, p. 1438], if L is a differential operator defined on the interval $(-\pi, \pi) = (-\pi, 0) \cup (0, \pi)$ and L_{\pm} are restrictions of L on $(-\pi, 0)$ and $(0, \pi)$, then $\sigma_c(L) = \sigma_c(L_+) \cup \sigma_c(L_-)$, where $\sigma_c(L)$ denotes the continuous spectrum of L . By the symmetry of the two intervals, it is sufficient to prove that the operator L_+ has no continuous spectrum on $(0, \pi)$ (independently of the boundary conditions at $\theta = 0$ and $\theta = \pi$). It is also sufficient to carry out the proof for $\epsilon > 0$. Let $f_+(t) = f(\theta)$ on $\theta \in [0, \pi]$ and

$$\cos \theta = \tanh t, \quad \sin \theta = \operatorname{sech} t, \quad t \in \mathbb{R},$$

such that the interval $[0, \pi]$ for θ is mapped to the infinite line \mathbb{R} for t . The function $f_+(t)$ satisfies the spectral problem

$$-\epsilon f_+''(t) + f_+'(t) = \lambda \operatorname{sech} t f_+(t). \tag{2.10}$$

With a transformation $f_+(t) = e^{t/2\epsilon} g_+(t)$, the spectral problem (2.10) is written in the symmetric form

$$-\epsilon g_+''(t) + \frac{1}{4\epsilon} g_+(t) = \lambda \operatorname{sech} t g_+(t). \tag{2.11}$$

Thus, our operator is extended to a symmetric operator with an exponentially decaying weight $\rho(t) = \operatorname{sech}(t)$. According to Corollary 3 in [8, p. 1437], if L is a symmetric operator on an open interval (a, b) and L_0 is a self-adjoint extension of L with respect to some boundary conditions at $x = a$ and $x = b$, then $\sigma_c(L) = \sigma_c(L_0)$. Here $a = -\infty$, $b = \infty$, and we need to show that the continuous spectrum of the symmetric problem (2.11) is empty in $L^2(\mathbb{R})$. This follows by Theorem 7 in [10, p. 93]: since the weight function $\rho(t)$ of the problem $-y''(t) - \lambda \rho(t)y(t) = 0$ on $t \in \mathbb{R}$ decays faster than $1/t^2$ as $|t| \rightarrow \infty$, the spectrum of $-y''(t) - \lambda \rho(t)y(t) = 0$ is purely discrete.¹ \square

3. Eigenvalues of the linear operator L

By results of Lemmas 2–5, the spectral problem (1.3) for $|\epsilon| < 2$ may have only two types of eigenvalues in addition to the simple zero eigenvalue: either pairs of purely imaginary eigenvalues or quartets of symmetric complex eigenvalues. We prove in Lemmas 6 and 7 that there exists an infinite sequence of eigenvalues λ which accumulate to infinity along the imaginary axis. Furthermore, we prove in Theorem 1 that the eigenfunctions associated to all eigenvalues of the spectral problem (1.3) form a complete dense set in $L^2_{\text{per}}([-\pi, \pi])$. In the end of this section, Theorem 2 gives a necessary and sufficient condition that the set of eigenfunctions forms a basis in $L^2_{\text{per}}([-\pi, \pi])$.

Lemma 6. *Let $0 < \epsilon < 2$ and $\epsilon \neq \frac{1}{n}$, $n \in \mathbb{N}$. For $\lambda \in \mathbb{C}$, the spectral problem (1.3) admits three sets of two linearly independent solutions in the form of the Frobenius series*

$$-\pi < \theta < \pi: \quad f_1 = 1 + \sum_{n \in \mathbb{N}} c_n \theta^n, \quad f_2 = \theta^{-1/\epsilon} \left(1 + \sum_{n \in \mathbb{N}} d_n \theta^n \right), \tag{3.1}$$

and

$$0 < \pm\theta < \pi: \quad f_1^{\pm} = 1 + \sum_{n \in \mathbb{N}} a_n^{\pm} (\pi \mp \theta)^n, \quad f_2^{\pm} = (\pi \mp \theta)^{1/\epsilon} \left(1 + \sum_{n \in \mathbb{N}} b_n^{\pm} (\pi \mp \theta)^n \right), \tag{3.2}$$

where all coefficients are uniquely defined. The solution $f_1(\theta)$ is an analytic function of $\lambda \in \mathbb{C}$ uniformly on $\theta \in [-\pi, \pi]$.

Proof. Existence of two linearly independent solutions on $-\pi < \theta < \pi$ in the form (3.1) and on $0 < \pm\theta < \pi$ in the form (3.2) follows by the Frobenius method² [15]. Since the spectral problem (1.3) depends analytically on λ and the Frobenius series converges absolutely and uniformly in between two singular points, the solution $f_1(\theta)$ is analytic in

¹ Although the spectral problem (2.11) has an additional term $Cy(t)$ with $C > 0$, this term only makes better the inequality (30) in the proof of Theorem 7 of [10, p. 93].

² The difference between the two indices of the indicial equation at $\theta = 0$ or $\theta = \pm\pi$ is $\frac{1}{\epsilon}$ and it is non-integer for $\epsilon \neq \frac{1}{n}$, $n \in \mathbb{N}$. An additional logarithmic term $\log(\pi - \theta)$ may be present in the Frobenius series if $\epsilon = \frac{1}{n}$, $n \in \mathbb{Z}$.

$\lambda \in \mathbb{C}$ for any fixed $\theta \in (-\pi, \pi)$. Due to uniqueness of the Frobenius series, the solution $f_1(\theta)$ can be equivalently represented by the other solutions

$$f_1(\theta) = A^\pm f_1^\pm(\theta) + B^\pm f_2^\pm(\theta), \quad 0 < \pm\theta < \pi, \tag{3.3}$$

where A^\pm and B^\pm are some constants, while the functions $f_1^\pm(\theta)$ and $f_2^\pm(\theta)$ are analytic in $\lambda \in \mathbb{C}$ for any fixed $\pm\theta \in (0, \pi]$. By matching analytic solutions for any $\pm\theta \in (0, \pi)$, we find that A^\pm and B^\pm are analytic functions of $\lambda \in \mathbb{C}$, the Frobenius series for $f_1(\theta)$ converges absolutely and uniformly on $[-\pi, \pi]$, and the solution $f_1(\theta)$ is an analytic function in $\lambda \in \mathbb{C}$. \square

Corollary 1. *There exists an analytic function $F_\epsilon(\lambda)$ on $\text{Im } \lambda > 0$, roots of which give isolated eigenvalues of the spectral problem (1.3) with the account of their multiplicity. The only accumulation point of isolated eigenvalues in the λ -plane may occur at infinity.*

Proof. The function $f \in H^1([-\pi, \pi])$ satisfies the spectral problem (1.3) if and only if $f(\theta) = C_0 f_1(\theta)$ on $\theta \in [-\pi, \pi]$, where $C_0 = 1$ thanks to the scaling invariance of homogeneous equations. By using the representation (3.3), we can find that $A^\pm = \lim_{\theta \rightarrow \pm\pi} f_1(\theta)$ are uniquely defined analytic functions in $\lambda \in \mathbb{C}$. The function $F_\epsilon(\lambda) = A^+ - A^-$ is analytic function of $\lambda \in \mathbb{C}$ by construction and zeros of $F_\epsilon(\lambda)$ on $\text{Im } \lambda > 0$ coincide with the eigenvalues λ of the spectral problem (1.3) with the account of their multiplicity. If $F_\epsilon(\lambda_0) = 0$ for some $\lambda_0 \in \mathbb{C}$, the corresponding eigenfunction $f(\theta)$ lies in $H^1_{\text{per}}([-\pi, \pi])$, i.e. it satisfies the periodic boundary conditions $f(\pi) = f(-\pi)$. By analytic function theory, the sequence of roots of $F_\epsilon(\lambda)$ cannot accumulate at a finite point on $\lambda \in \mathbb{C}$. \square

Remark 2. We will use the method involving the analytic function $F_\epsilon(\lambda)$ on $\lambda \in \mathbb{C}$ for a numerical shooting method which enables us to approximate eigenvalues of the spectral problem (1.3). This method involves less computations than the shooting method described in Appendix C of [3].

Lemma 7. *Fix $0 < \epsilon < 2$ and let $\{\lambda_n\}_{n \in \mathbb{N}}$ be a set of eigenvalues of the spectral problem (1.3) with $\text{Im } \lambda_n > 0$, ordered in the ascending order of $|\lambda_n|$. There exists a finite number $N \geq 1$, such that for all $n \geq N$, $\lambda_n = i\omega_n \in i\mathbb{R}_+$ and*

$$\omega_n = Cn^2 + o(n^2) \quad \text{as } n \rightarrow \infty, \tag{3.4}$$

for some $C > 0$.

Proof. We reduce the spectral problem (1.3) to two uncoupled Schrödinger equations on an infinite line. Let $f(\theta)$ be represented on two intervals $\pm\theta \in [0, \pi]$ by using the transformations

$$\cos \theta = \tanh t, \quad \sin \theta = \pm \text{sech } t, \tag{3.5}$$

where $t \in \mathbb{R}$. Then, the functions $f_\pm(t) = f(\theta)$ on $\pm\theta \in [0, \pi]$ satisfy the uncoupled spectral problems

$$-\epsilon f_\pm''(t) + f_\pm'(t) = \pm \lambda \text{sech } t f_\pm(t), \quad t \in \mathbb{R}. \tag{3.6}$$

The normalization condition $f(0) = 1$ is equivalent to the condition $\lim_{t \rightarrow \infty} f_\pm(t) = 1$. The periodic boundary condition $f(\pi) = f(-\pi)$ is equivalent to the condition $\lim_{t \rightarrow -\infty} f_-(t) = \lim_{t \rightarrow -\infty} f_+(t)$. The linear problems (3.6) are reformulated as the quadratic Riccati equations by using the new variables

$$f_\pm(t) = e^{\int_\infty^t S_\pm(t') dt'}: \quad S_\pm - \epsilon(S'_\pm + S_\pm^2) = \pm \lambda \text{sech } t. \tag{3.7}$$

We choose a negative root of the quadratic equation in the form

$$S_\pm(t) = \frac{1 - \sqrt{1 \mp 4\epsilon \lambda \text{sech } t - 4\epsilon^2 R_\pm}}{2\epsilon}, \quad R_\pm = S'_\pm(t). \tag{3.8}$$

The representation (3.8) becomes the chain fraction if the derivative of $S_\pm(t)$ is defined recursively from the same expression (3.8). By using the theory of chain fractions, we claim that $R_\pm = O(\sqrt{|\lambda|})$ as $|\lambda| \rightarrow \infty$ uniformly on $t \in \mathbb{R}$. The function $F_\epsilon(\lambda)$ of Corollary 1 is now expressed by

$$F_\epsilon(\lambda) = \lim_{t \rightarrow -\infty} [f_+(t) - f_-(t)] = e^{\int_{-\infty}^{\infty} S_+(t) dt} - e^{\int_{-\infty}^{\infty} S_-(t) dt}. \tag{3.9}$$

Zeros of $F_\epsilon(\lambda)$ are equivalent to zeros of the infinite set of functions

$$G_n(\lambda) = \frac{1}{4\pi i \epsilon} \int_{-\infty}^{\infty} [\sqrt{1 + 4\epsilon\lambda \operatorname{sech} t - 4\epsilon^2 R_-(t)} - \sqrt{1 - 4\epsilon\lambda \operatorname{sech} t - 4\epsilon^2 R_+(t)}] dt - n, \tag{3.10}$$

where $n \in \mathbb{N}$. If $R_\pm(t) \equiv 0$, the function $\tilde{G}_n(\omega) = G_n(i\omega)$, $n \in \mathbb{N}$ is real-valued and strictly increasing on $\omega \in \mathbb{R}_+$ with $\tilde{G}_n(0) = -n$. By performing asymptotic analysis, we compute that

$$\begin{aligned} & \frac{1}{4\pi i \epsilon} \int_{-\infty}^{\infty} [\sqrt{1 + 4i\epsilon\omega \operatorname{sech} t - 4\epsilon^2 R_-(t)} - \sqrt{1 - 4i\epsilon\omega \operatorname{sech} t - 4\epsilon^2 R_+(t)}] dt \\ &= \frac{1}{\pi i} \int_{-\infty}^{\infty} \frac{2i\omega \operatorname{sech} t + \epsilon(R_+ - R_-)}{\sqrt{1 + 4i\epsilon\omega \operatorname{sech} t - 4\epsilon^2 R_-(t)} + \sqrt{1 - 4i\epsilon\omega \operatorname{sech} t - 4\epsilon^2 R_+(t)}} dt \\ &= \frac{\sqrt{\omega}}{\sqrt{2\epsilon\pi}} \int_{-\infty}^{\infty} \frac{dt}{\sqrt{\cosh t}} + o(\sqrt{\omega}), \end{aligned} \tag{3.11}$$

such that $\lim_{\omega \rightarrow \infty} \tilde{G}_n(\omega) = \infty$. Therefore, there exists exactly one root $\omega = \omega_n$ of $\tilde{G}_n(\omega)$ for each n . Since $R_- = \bar{R}_+$ for $\lambda = i\omega \in i\mathbb{R}$, each simple root of $\tilde{G}_n(\omega)$ persists for non-zero values of $R_\pm(t) = O(\sqrt{\omega})$ uniformly on $t \in \mathbb{R}$ as $\omega \rightarrow \infty$. According to the asymptotic result (3.11), the roots ω_n of $\tilde{G}_n(\omega)$ satisfy the asymptotic distribution (3.4) with $C = \frac{2\epsilon\pi^2}{(\int_{-\infty}^{\infty} \frac{dt}{\sqrt{\cosh t}})^2}$. \square

Remark 3. Analysis of Lemma 7 extends the formal WKB approach proposed in Section 3 of [3]. In particular, Eq. (3.10) with $R_\pm = 0$ has been obtained in Eq. (3.11) of [3].

Definition 1. A Schauder basis for a Banach space X is a sequence $\{f_n\}_{n \in \mathbb{N}}$ in X with the property that every f in X has a unique representation of the form $f = \sum_{n=1}^{\infty} c_n f_n$ with some coefficients $\{c_n\}_{n \in \mathbb{N}}$, in which the N th partial sum is convergent in the X -norm.

Definition 2. The set of functions $\{f_n\}_{n \in \mathbb{N}}$ is said to be complete in a Banach space X if any function f in X can be approximated by a finite linear combination $f_N = \sum_{n=1}^N c_n f_n$ in the following sense: for any fixed $\epsilon > 0$, there exists $N \geq 1$ and a certain set of coefficients $\{c_1, c_2, \dots, c_N\}$, such that the inequality $\|f - f_N\|_X < \epsilon$ holds. The complete set of functions $\{f_n\}_{n \in \mathbb{N}}$ is said to be minimal if it is linear independent in the sense of finite linear combinations.

Theorem 1. Let $\{f_n\}_{n \in \mathbb{N}}$ be the set of eigenfunctions corresponding to the set of eigenvalues $\{\lambda_n\}_{n \in \mathbb{N}}$ in Lemma 7 with $\operatorname{Im} \lambda_n > 0$. The set of eigenfunctions is complete in the Banach space

$$X_0 = \left\{ f \in L^2_{\text{per}}([- \pi, \pi], \mathbb{C}) : \int_{-\pi}^{\pi} f(\theta) d\theta = 0 \right\}.$$

Proof. By Corollary 1, eigenvalues of L with $\operatorname{Im} \lambda > 0$ accumulate to infinity, such that the operator $M = L^{-1}$ acting on elements in X_0 is compact. By Lemma 7, there are infinitely many eigenvalues of L and large eigenvalues are all purely imaginary, such that $|\lambda_n| = O(n^2)$ as $n \rightarrow \infty$. These two facts can be used to show that two sufficient conditions of the Lidskii’s Completeness Theorem are satisfied. According to Theorem 6.1 in [11, p. 302], the set of eigenvectors and generalized eigenvectors of a compact operator M in a Hilbert space X_0 is complete if there exists $p > 0$ such that

$$s_n(M) = o(n^{-\frac{1}{p}}) \quad \text{as } n \rightarrow \infty, \tag{3.12}$$

where s_n is a singular number of the operator M , and the set

$$W_M = \{(Mf, f) : f \in X_0, \|f\|_{X_0} = 1\} \tag{3.13}$$

lies in a closed angle θ_M with vertex at 0 and opening $\frac{\pi}{p}$. Since the singular numbers s_n are eigenvalues of the positive self-adjoint operator $(MM^*)^{1/2}$ and the eigenvalues of L grow like $O(n^2)$ as $n \rightarrow \infty$, we have $s_n(M) = O(n^{-2})$ as $n \rightarrow \infty$, such that the first condition (3.12) is verified with $p = 1$. Since all $\text{Im } \lambda_n > 0$ for the set of eigenvalues $\{\lambda_n\}_{n \in \mathbb{N}}$ of Lemma 7, the spectrum of M lies in the lower half plane, such that the second condition (3.13) is also verified with $p = 1$ ($\theta_M = \pi$). \square

Corollary 2. *The set of eigenfunctions $\{f_n\}_{n \in \mathbb{Z}}$ with $f_0 = 1$ and $f_{-n} = \bar{f}_n, \forall n \in \mathbb{N}$ is complete in $L^2_{\text{per}}([-\pi, \pi])$.*

Remark 4. Due to linear independence of eigenfunctions for distinct eigenvalues, the complete set of eigenfunctions $\{f_n\}_{n \in \mathbb{Z}}$ is also minimal if all eigenvalues are simple.³

Theorem 2. *Let $\{f_n\}_{n \in \mathbb{Z}}$ be a complete and minimal set of eigenfunctions of the spectral problem (1.3) for the set of eigenvalues $\{\lambda_n\}_{n \in \mathbb{Z}}$ in Theorem 1. The set of eigenfunctions forms a basis in Hilbert space $L^2_{\text{per}}([-\pi, \pi])$ if and only if $\lim_{n \rightarrow \infty} \cos(\widehat{f_n, f_{n+1}}) < 1$.*

Proof. According to Theorem 2 in [14, p. 31], the complete and minimal set of eigenfunctions $\{f_n\}_{n \in \mathbb{Z}}$ forms a basis in Hilbert space $X = L^2_{\text{per}}([-\pi, \pi])$ if and only if $\sup_N \|P_N\| < \infty$, where P_N is a projector from X to a subspace spanned by $\{f_n\}_{-N \leq n \leq N}$ parallel to the subspace spanned by $\{f_n\}_{|n| > N+1}$. Since the Hilbert space X is a direct sum of the two subspaces above, the norm of the parallel projector P_N has the geometrical representation $\|P_N\| = \frac{1}{\sin \alpha_N}$, where α_N is the angle between the two subspaces [1]. This implies that the set $\{f_n\}_{n \in \mathbb{Z}}$ is a basis in the Hilbert space X if and only if

$$\cos(\widehat{f_n, f_{n+1}}) = \frac{|(f_n, f_{n+1})|}{\|f_n\| \|f_{n+1}\|} < 1, \tag{3.14}$$

for sufficiently large $n \in \mathbb{Z}$ [12]. \square

4. Numerical approximations

We approximate isolated eigenvalues of the spectral problem (1.3) for $0 < \epsilon < 2$ numerically. In agreement with numerical results in [3], we show that all eigenvalues in the set $\{\lambda_n\}_{n \in \mathbb{Z}}$ are simple and purely imaginary. Therefore, the set $\{\lambda_n\}_{n \in \mathbb{Z}}$ can be ordered in the ascending order, such that $\lambda_0 = 0, \lambda_n = -\lambda_{-n}, \forall n \in \mathbb{N}, \text{Im } \lambda_n < \text{Im } \lambda_{n+1}$ and $\lim_{n \rightarrow \infty} |\lambda_n| = \infty$. We also show that the angle between two subsequent eigenfunctions $f_n(\theta)$ and $f_{n+1}(\theta)$ in the set $\{f_n(\theta)\}_{n \in \mathbb{Z}}$ tends to zero as $n \rightarrow \infty$. All computations of this section are performed in MATLAB 6 under the Windows platform.

4.1. Shooting method

The numerical shooting method is based on the ODE formulation of the spectral problem (1.3). By Lemma 6 and Corollary 1, complex eigenvalues $\lambda \in \mathbb{C}$ are determined by roots of the analytic function $F_\epsilon(\lambda)$ in the λ -plane. The number of complex eigenvalues can be computed with the winding number theory. The number and location of purely imaginary eigenvalues can be found from real-valued roots of a scalar real-valued function.

Proposition 1. *Let the eigenfunction $f(\theta)$ of the spectral problem (1.3) for $0 < \epsilon < 2$ be normalized by the condition $f(0) = 1$. The eigenvalue λ is purely imaginary if and only if $f(\theta) = \bar{f}(-\theta)$ on $\theta \in [-\pi, \pi]$.*

³ By Lemma 7, all eigenvalues are simple starting with some $n \geq N$.

Proof. If $\lambda \in i\mathbb{R}$ and $f(\theta)$ satisfies the second-order ODE (1.3) on $\theta \in [-\pi, \pi]$, then $\bar{f}(-\theta)$ satisfies the same ODE (1.3) on $\theta \in [-\pi, \pi]$. By Corollary 1, if $f \in H^1_{\text{per}}([-\pi, \pi])$, $f(0) = 1$ and $0 < \epsilon < 2$, the solution $f(\theta)$ is uniquely defined. By uniqueness of solutions, $f(\theta) = \bar{f}(-\theta)$ on $\theta \in [-\pi, \pi]$.

If $f(\theta) = \bar{f}(-\theta)$ on $\theta \in [-\pi, \pi]$, then

$$\int_{-\pi}^{\pi} \sin \theta |f'(\theta)|^2 d\theta = \int_0^{\pi} \sin \theta |f'(\theta)|^2 d\theta - \int_0^{\pi} \sin \theta |f'(-\theta)|^2 d\theta = 0,$$

such that $\text{Re } \lambda = 0$ according to the equality (2.6) in Lemma 3. \square

Corollary 3. Let $f(\theta)$ be an eigenfunction of the spectral problem (1.3) for $\lambda \in i\mathbb{R}$, such that $f \in H^1_{\text{per}}([-\pi, \pi])$ and $f(0) = 1$. Then, $f(\pi) = f(-\pi)$ is equivalent to $f(\pi) \in \mathbb{R}$. The eigenvalue $\lambda \in i\mathbb{R}$ is simple if and only if

$$(f^*, f) = 2 \text{Re} \int_0^{\pi} f(\theta) \bar{f}(\pi - \theta) d\theta \neq 0, \tag{4.1}$$

where $f^* = f(\pi - \theta)$ is an eigenfunction of the adjoint operator L^* for the same eigenvalue.

Proof. The first assertion follows by the symmetry relation $f(\theta) = \bar{f}(-\theta)$ evaluated at $\theta = \pi$. The second assertion follows by Lemma 2 with the use of the symmetry $f^*(\theta) = f(\pi - \theta)$. \square

By Lemma 6, the function $f(\theta)$ with $f(0) = 1$ is represented uniquely by the Frobenius series

$$f(\theta) = f_1(\theta) = 1 + \sum_{n \in \mathbb{N}} c_n \theta^n, \tag{4.2}$$

where the coefficients $\{c_n\}_{n \in \mathbb{N}}$ are uniquely defined by the recursion relation

$$c_n = -\frac{1}{n(1 + \epsilon n)} \left(\lambda c_{n-1} + \epsilon n \sum_{m \in [1, n-2]'} \frac{(-1)^{\frac{n-m}{2}} m}{(n-m+1)!} c_m \right), \quad n \in \mathbb{N}, \tag{4.3}$$

where $c_0 = 1$ and $[1, n-2]'$ is a set of integers in the interval $[1, n-2]$ such that $n-m$ is even. For instance,

$$c_1 = -\frac{\lambda}{1 + \epsilon}, \quad c_2 = \frac{\lambda^2}{2(1 + \epsilon)(1 + 2\epsilon)}, \quad c_3 = -\frac{\lambda(\lambda^2 + \epsilon(1 + 2\epsilon))}{3!(1 + \epsilon)(1 + 2\epsilon)(1 + 3\epsilon)}$$

and so on. We truncate the power series expansion on $N = 100$ terms and approximate the initial value $[f(\theta_0), f'(\theta_0)]$ at $\theta_0 = 10^{-8}$. By using the fourth-order Runge–Kutta ODE solver with time step $h = 10^{-4}$, we obtain a numerical approximation of $f \equiv f_+(\theta)$ on $\theta \in [\theta_0, \pi - \theta_0]$ for λ and $f \equiv f_-(\theta)$ on the same interval for $-\lambda$. By Lemma 2(i), the numerical approximation of the function $F_\epsilon(\lambda)$ of Corollary 1 is

$$\widehat{F}_\epsilon(\lambda) = f_+(\pi - \theta_0) - f_-(\pi - \theta_0). \tag{4.4}$$

If $\lambda \in i\mathbb{R}$, the function $\widehat{F}_\epsilon(\lambda)$ is simplified by using Corollary 3 as $\widehat{F}_\epsilon(\lambda) = 2i \text{Im } f_+(\pi - \theta_0)$. Table 1 represents the numerical approximations of the first four non-zero eigenvalues $\lambda \in i\mathbb{R}$ for $\epsilon = 0.5, 1.0, 1.5^4$ with the error computed from the residual

$$R = \left| \frac{(f, Lf)}{(f, f)} - \lambda \right|.$$

We can see from Table 1 that the accuracy drops with larger values of ϵ and for larger eigenvalues, but the eigenvalues persist inside the interval $|\epsilon| < 2$.

⁴ We note that the Frobenius series (4.2) is not affected by the logarithmic terms for $\epsilon = 0.5$ and $\epsilon = 1.0$, since 0 is the largest index of the indicial equation at $\theta = 0$.

Table 1

Numerical approximations of the first four eigenvalues $\lambda = i\omega_n$ of the spectral problem (1.3) and the residuals $R = R_n$ for three values of ϵ

ϵ	ω_1	R_1	ω_2	R_2	ω_3	R_3	ω_4	R_4
0.5	1.167342	0.000051	2.968852	0.000405	5.483680	0.001436	8.715534	0.003653
1.0	1.449323	0.000837	4.319645	0.007069	8.631474	0.024964	14.382886	0.061881
1.5	1.757278	0.002691	5.719671	0.018412	11.846709	0.054271	20.138824	0.113834




Fig. 1. The real part (dark) and imaginary part (bright) of the eigenfunction $f(\theta)$ on $\theta \in [0, \pi]$ for the first (solid) and second (dashed) eigenvalues $\lambda = i\omega_{1,2} \in i\mathbb{R}_+$ for $\epsilon = 0.5$ (left) and $\epsilon = 1.5$ (right).




Fig. 2. The image of the curve $w = \widehat{F}_\epsilon(\lambda)$, when λ traverses along the contours Λ_1, Λ_2 and Λ_3 for $\epsilon = 0.5$: the image curve on the w -plane (left) and the argument of w (right).

Fig. 1 shows the profiles of eigenfunctions $f(\theta)$ on $\theta \in [0, \pi]$ for the first two eigenvalues $\lambda = i\omega_{1,2} \in i\mathbb{R}_+$ for $\epsilon = 0.5$ (left) and $\epsilon = 1.5$ (right). We can see from Fig. 1 that the derivative of $f(\theta)$ becomes singular as $\theta \rightarrow \pi^-$ for $\epsilon \geq 1$. We can also see that the real part of the eigenfunction $f(\theta)$ has one zero on $\theta \in (0, \pi)$ for the first eigenvalue and two zeros for the second eigenvalue, while the imaginary part of the eigenfunction $f(\theta)$ has one less number of zeros. The numerical approximations of the eigenvalue and eigenfunctions of the spectral problem (1.3) are structurally stable with respect to variations in θ_0, N and h .

Fig. 2 shows the complex plane of $w = \widehat{F}_\epsilon(\lambda)$ (left) and the argument of w (right) when λ traverses along the first quadrant of the complex plane $\lambda \in \Lambda_1 \cup \Lambda_2 \cup \Lambda_3$ for $\epsilon = 0.5$. Here $\Lambda_1 = x + ir$ with $x \in [r, R]$, $\Lambda_2 = Re^{i\varphi}$ with $\varphi \in [\varphi_0, \frac{\pi}{2} - \varphi_0]$ and $\Lambda_3 = r + iy$ with $y \in [r, R]$, where $r = 0.1, R = 10$, and $\varphi_0 = \arctan(r/R)$. It is obvious that the

winding number of $\widehat{F}_\epsilon(\lambda)$ across the closed contour is zero. Therefore, no zeros of $\widehat{F}_\epsilon(\lambda)$ occurs in the first quadrant of the complex plane $\lambda \in \mathbb{C}$. The numerical result is structurally stable with respect to variations in r , R and ϵ .

4.2. Spectral method

The numerical spectral method is based on the reformulation of the second-order ODE (1.3) as the second-order difference equation and the subsequent truncation of the difference eigenvalue problem. It is found in [17] that the truncation procedure lead to spurious complex eigenvalues which bifurcate off the imaginary axis.

Let $f \in H^1_{\text{per}}([-\pi, \pi])$ be an eigenfunction of the spectral problem (1.3). This eigenfunction is equivalently represented by the Fourier series

$$f(\theta) = \sum_{n \in \mathbb{Z}} f_n e^{-in\theta}, \quad f_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(\theta) e^{in\theta} d\theta, \tag{4.5}$$

where the infinite-dimensional vector $\mathbf{f} = (\dots, f_{-2}, f_{-1}, f_0, f_1, f_2, \dots)$ is defined in $\mathbf{f} \in l^2_1(\mathbb{Z})$ equipped with the norm $\|\mathbf{f}\|_{l^2_1}^2 = \sum_{n \in \mathbb{Z}} (1 + n^2) |f_n|^2 < \infty$. The spectral problem (1.3) for $|\epsilon| < 2$ is equivalent to the difference eigenvalue problem

$$n f_n + \frac{\epsilon}{2} n [(n + 1) f_{n+1} - (n - 1) f_{n-1}] = -i \lambda f_n, \quad n \in \mathbb{Z}. \tag{4.6}$$

The difference eigenvalue problem (4.6) splits into three parts

$$A \mathbf{f}_+ = -i \lambda \mathbf{f}_+, \quad A \mathbf{f}_- = i \lambda \mathbf{f}_-, \quad \lambda f_0 = 0, \tag{4.7}$$

where $\mathbf{f}_\pm = (f_{\pm 1}, f_{\pm 2}, \dots)$ and A is an infinite-dimensional matrix

$$A = \begin{bmatrix} 1 & \epsilon & 0 & 0 & \dots \\ -\epsilon & 2 & 3\epsilon & 0 & \dots \\ 0 & -3\epsilon & 3 & 6\epsilon & \dots \\ 0 & 0 & -6\epsilon & 4 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}. \tag{4.8}$$

Since $A = D - iS$, where D is a diagonal matrix and S is a self-adjoint tri-diagonal matrix, one can define the discrete counterpart of Lemma 3

$$\text{Im } \lambda = \frac{(\mathbf{f}_+, D \mathbf{f}_+)}{(\mathbf{f}_+, \mathbf{f}_+)} = \frac{\sum_{n \in \mathbb{N}} n |f_n|^2}{\sum_{n \in \mathbb{N}} |f_n|^2}, \quad \text{Re } \lambda = \frac{(\mathbf{f}_+, S \mathbf{f}_+)}{(\mathbf{f}_+, \mathbf{f}_+)},$$

where $\text{Im } \lambda > 0$. The adjoint eigenfunction $f^*(\theta) = f(\pi - \theta)$ is recovered from the eigenvector \mathbf{f} by $\mathbf{f}^* = J \mathbf{f}$, where

$$J = \begin{bmatrix} 0 & 0 & J_0 \\ 0 & 1 & 0 \\ J_0 & 0 & 0 \end{bmatrix}$$

and J_0 is a diagonal operator with entries $(-1, 1, -1, 1, \dots)$.

According to Theorem 1, rewritten from the set of eigenfunctions $\{f_n\}_{n \in \mathbb{Z}}$ to the set of eigenvectors $\{\mathbf{f}_n\}_{n \in \mathbb{Z}}$, the inverse matrix operator A^{-1} is of the Hilbert–Schmidt type, and hence compact. Let $A_N^{-1} = P_N A^{-1} P_N$ denote the truncation of the matrix operator A^{-1} at the first N rows and columns, where P_N is an orthogonal projector from an infinite-dimensional vector to the N -dimensional vector of the first N components.

Proposition 2. *Operator sequence A_N^{-1} converges uniformly to the compact operator A^{-1} as $N \rightarrow \infty$. Eigenvalues of the matrices A_N^{-1} converge to the eigenvalues of the compact operator A^{-1} as $N \rightarrow \infty$.*

Proof. It follows from the Finite Rank Approximation Theorem that $P_N A^{-1}$ converges uniformly to the compact operator A^{-1} . Therefore, for any $\epsilon > 0$, there exists a number $N_1 \geq 1$ such that

$$\forall N > N_1: \quad \|P_N A^{-1} - A^{-1}\| < \frac{\epsilon}{2}.$$

Because the adjoint operator is also compact and the orthogonal projector P_N is a self-adjoint operator, the sequence $P_N^* A^{-1*}$ uniformly converges to A^{-1*} . Therefore, for any $\epsilon > 0$, there exists a number $N_2 \geq 1$ such that

$$\forall N > N_2: \quad \|P_N^* A^{-1*} - A^{-1*}\| < \frac{\epsilon}{2}.$$

Let $N_0 = \max(N_1, N_2)$. For any $N > N_0$, we have

$$\begin{aligned} \|A^{-1} - P_N A^{-1} P_N\| &= \|(A^{-1} - P_N A^{-1}) + P_N(A^{-1*} - P_N^* A^{-1*})^*\| \\ &\leq \|A^{-1} - P_N A^{-1}\| + \|P_N\| \|(A^{-1*} - P_N^* A^{-1*})^*\| \\ &\leq \|A^{-1} - P_N A^{-1}\| + \|A^{-1*} - P_N^* A^{-1*}\| \leq \epsilon. \end{aligned}$$

Therefore, $\lim_{N \rightarrow \infty} A_N^{-1} = A^{-1}$.

Let $\lambda_0 \neq 0$ belongs to the spectrum of the operator A^{-1} . Because all eigenvalues are isolated, there exists an open ball $D_0 \in \text{Dom}(A^{-1})$ with the boundary ∂D_0 passing through regular points of operator A such that λ_0 is the only point of D_0 in the spectrum set of A^{-1} . It follows from the compactness of ∂D_0 that the set $\{(A_N^{-1} - \lambda I)^{-1}: \lambda \in \partial D_0\}$ is uniformly bounded by N and by λ . Therefore, the sequence of the Riesz projectors

$$R_N = -\frac{1}{2\pi i} \oint_{\Gamma_{D_0}} (A_N^{-1} - \lambda I)^{-1} d\lambda$$

strongly converges to the limiting projector

$$R = -\frac{1}{2\pi i} \oint_{\Gamma_{D_0}} (A^{-1} - \lambda I)^{-1} d\lambda.$$

If all $R_N = 0$, then the limiting projector $R = 0$. \square

Remark 5. The distance between eigenvalues of A_N^{-1} and A^{-1} may not be small for fixed N , but for every fixed eigenvalue it becomes small for large N . The convergence of eigenvalues is not uniform in N .

The smallest eigenvalues of the truncated matrix A_N^{-1} are found with the parallel Krylov subspace iteration algorithm [9]. Fig. 3 shows the distance between eigenvalues of the shooting method and eigenvalues of the Krylov spectral method for $\epsilon = 0.1$. The difference between two eigenvalues is small of the order $O(10^{-3})$ but the advantage of the parallel algorithm is that the calculating time of 20 largest eigenvalues of A_N^{-1} for $N = 10^6$ takes less than one minute on a network of 16 processors while finding the same set of eigenvalues by the shooting method with the time step $h = 10^{-5}$ takes about one hour.

Fig. 4 shows symmetric pairs of eigenvalues of the matrix A_N for $\epsilon = 0.3$ at $N = 128$ (left) and $N = 1024$ (right). We confirm the numerical result of [17] that the truncation of the matrix operator A always produces splitting of large eigenvalues off the imaginary axis.⁵ However, the larger is N , the more eigenvalues remain on the purely imaginary axis. Therefore, the corresponding eigenvectors can be used to compute the angle in Theorem 2.

Fig. 5 (left) show the values of the cosine of the angle (3.14) for the first 20 purely imaginary eigenvalues for $\epsilon = 0.1$. As we can see from the figure, the angle between two eigenvectors tends to zero for larger eigenvalues up to the numerical accuracy. Fig. 5 (right) and Table 2 show that the angle between the first two eigenvectors drops to zero faster with larger values of the parameter ϵ .

⁵ A similar phenomenon called the spectral pollution has been investigated in [7]. If the variational minimum–maximum principle is not applicable to a non-self-adjoint problem, as in our case, eigenvalues of the spectral method may differ drastically from eigenvalues of the continuous problem.




Fig. 3. The distance between eigenvalues computed by the shooting and spectral methods for $\epsilon = 0.1$.




Fig. 4. Spectrum of the truncated difference eigenvalue problem (4.6) for $\epsilon = 0.3$: $N = 128$ (left) and $N = 1024$ (right).




Fig. 5. Left: the values of $\cos(\widehat{f_n, f_{n+1}})$ for the first 20 purely imaginary eigenvalues for $\epsilon = 0.1$. Right: the values of $\cos(\widehat{f_1, f_2})$ versus ϵ .

Table 2
Numerical values of $\cos(\widehat{f_n, f_{n+1}})$ for the first 16 purely imaginary eigenvalues for three values of ϵ

Eigenvectors	$\epsilon = 0.1$	$\epsilon = 0.3$	$\epsilon = 0.5$
1–2	0.120166	0.325116	0.431987
2–3	0.461330	0.716192	0.780641
3–4	0.680709	0.838889	0.878055
4–5	0.799235	0.890440	0.914622
5–6	0.858944	0.921498	0.940306
6–7	0.892869	0.940395	0.955239
7–8	0.914745	0.953124	0.965235
8–9	0.930023	0.962120	0.972204
9–10	0.941262	0.968732	0.977265
10–11	0.949843	0.973741	0.981057
11–12	0.956580	0.977629	0.983988
12–13	0.961987	0.980702	0.986072
13–14	0.966407	0.983297	0.989617
14–15	0.970073	0.983459	0.990547
15–16	0.973153	0.995335	0.999101
16–17	0.975764	0.998749	0.999601




Fig. 6. The condition number for the first 40 purely imaginary eigenvalues for $\epsilon = 0.001$ (bright) and $\epsilon = 0.002$ (dark).

The angle between two subsequent eigenvectors is closely related to the condition number [16]

$$\text{cond}(\lambda_n) = \frac{\|f_n\| \|f_n^*\|}{|(f_n, f_n^*)|}. \tag{4.9}$$

By Lemma 2(iii), the condition number is infinite for multiple eigenvalues since $(f_n, f_n^*) = 0$. From the point of numerical accuracy, the larger is the condition number, the poorer is the structural stability of the numerically obtained eigenvalues to the truncation and round-off errors.

Fig. 6 shows the condition number (4.9) computed for the first 40 purely imaginary eigenvalues for $\epsilon = 0.001$ and $\epsilon = 0.002$. We can see that the condition number grows for larger eigenvalues which indicate their structural instability. Indeed, starting with some number n , all eigenvalues are no longer purely imaginary, according to the numerical approximations in Fig. 4. The condition number becomes extremely large with larger values of ϵ .

We finally illustrate that all true eigenvalues of the spectral problem (1.3) are purely imaginary and simple. To do so, we construct numerically the sign-definite imaginary type function and obtain the interlacing property of eigenvalues of the spectral problem (1.3) for two values $\epsilon = \epsilon_0$ and $\epsilon = \epsilon_1$, where $|\epsilon_1 - \epsilon_0|$ is small. We say that the eigenvalues exhibit the interlacing property if there exists an eigenvalue for $\epsilon = \epsilon_1$ between each pair of eigenvalues for $\epsilon = \epsilon_0$ and vice versa.

Table 3
The interlacing property of the first 15 purely imaginary eigenvalues for $\epsilon = 0.48$ and $\epsilon = 0.5$

$\text{Im } \lambda_{\epsilon_0}$	R_{ϵ_0}	$\text{Im } \lambda_{\epsilon_1}$	R_{ϵ_1}
1.063112	2.3244e-10	1.068314	2.4073e-10
2.970880	2.1967e-10	3.024428	2.2531e-10
5.414789	2.2024e-10	5.542829	2.2683e-10
8.471510	2.0904e-10	8.693066	2.1572e-10
12.312548	2.0079e-10	12.665485	2.0601e-10
16.816692	1.9765e-10	17.327038	2.0288e-10
22.014084	1.9617e-10	22.711070	2.0197e-10
27.899896	1.9527e-10	28.812177	2.0157e-10
34.474785	1.9501e-10	35.631088	2.0190e-10
41.738699	1.9558e-10	43.167733	2.0313e-10
49.691673	1.9671e-10	51.422281	2.0476e-10
58.333258	1.9796e-10	60.391382	2.0623e-10
67.665387	1.9904e-10	70.140636	2.0725e-10
77.957871	1.9989e-10	79.828287	2.0782e-10
89.484519	2.6566e-10	91.544035	2.0821e-10

A meromorphic function $G(\lambda)$ is called a sign-definite imaginary type function if $\text{Im } G(\lambda) \leq 0$ ($\text{Im } G(\lambda) \geq 0$) on $\text{Im}(\lambda) \leq 0$ ($\text{Im}(\lambda) \geq 0$) [2]. We construct the meromorphic function $G(\omega)$ in the form $G(\lambda) = \frac{F_{\epsilon_0}(\lambda)}{F_{\epsilon_1}(\lambda)}$, where $F_{\epsilon}(\lambda)$ is an analytical function of Corollary 1. The numerical approximation of the meromorphic function $G(\lambda)$ is given by $\widehat{G}(\lambda) = \frac{\widehat{F}_{\epsilon_0}(\lambda)}{\widehat{F}_{\epsilon_1}(\lambda)}$. According to Theorems II.2.1–II.3.1 in [2, pp. 437–439], the function $\widehat{G}(\lambda)$ is a meromorphic function of sign-definite imaginary type if and only if it has the form $\widehat{G}(\lambda) = \frac{P(\lambda)}{Q(\lambda)}$ where $P(\lambda)$ and $Q(\lambda)$ are polynomials with real coefficients, with real and simple zeros, which are interlacing.

Table 3 shows this interlacing property of eigenvalues for $\epsilon_0 = 0.48$ and $\epsilon_1 = 0.5$. The remainder term $R_{\epsilon} = \frac{\|L_f - \lambda f\|}{\|\lambda f\|}$ measures the numerical error of computations. We have also computed numerically the values of $\widehat{G}(\lambda)$ on the grid $0.1 < \text{Im } \lambda < 100$ and $0.1 < \text{Re } \lambda < 100$ with step size 0.1 in both directions (not shown). Based on the numerical data, we have confirmed that the function $\widehat{G}(\lambda)$ does indeed belongs to the class of sign-definite imaginary type functions while the eigenvalues $\{\lambda_n\}_{n \in \mathbb{Z}}$ exhibit the interlacing property. This computation gives a numerical verification that all eigenvalues of the spectral problem (1.3) are simple and purely imaginary.

5. Cauchy problem for the heat equation

We have proved that the operator L associated with the heat equation (1.1) admits a closure in $L^2_{\text{per}}([-\pi, \pi])$ with a domain in $H^1_{\text{per}}([-\pi, \pi])$ for $|\epsilon| < 2$. The spectrum of L consists of eigenvalues of finite multiplicities. Using the analytic function theory and the Fourier series, we have approximated eigenvalues numerically and showed that all eigenvalues of the spectral problem (1.3) are purely imaginary. Furthermore, we have proved with the assistance of numerical computations that the set of eigenfunctions of the spectral problem (1.3) is complete but does not form a basis in the Hilbert space $L^2_{\text{per}}([-\pi, \pi])$.

We shall show that these properties of the linear operator L are related to ill-posedness of the Cauchy problem for the periodic heat equation (1.1). According to the Hille–Yosida Theorem (see Section IX.7 in [18]), if L is a linear operator with a dense domain in a Banach space X and the resolvent operator $(I - \lambda^{-1}L)^{-1}$ exists for any $\text{Re } \lambda > 0$, then L is the infinitesimal generator of a strongly continuous semigroup if and only if

$$\|(I - \lambda^{-1}L)^{-1}\|_{X \rightarrow X} \leq C, \tag{5.1}$$

for some $C > 0$ uniformly in $\text{Re } \lambda > 0$. Moreover, if $C \leq 1$, then the semi-group is a contraction. The Cauchy problem associated with the operator L is well-posed if the conditions of the Hille–Yosida Theorem are satisfied and it is ill-posed otherwise.

According to the numerical results on pseudo-spectra in [3] and [17], the level set of the resolvent norm

$$R(\lambda) = \|(\lambda I - L)^{-1} \|_{L^2_{\text{per}}([-\pi, \pi]) \rightarrow L^2_{\text{per}}([-\pi, \pi])}$$

extends to the right half-plane, such that $R(\lambda)$ does not decay along the level set curves with $\text{Re } \lambda > 0$. This numerical fact serves as an indication that the conditions of the Hille–Yosida Theorem are not satisfied and the Cauchy problem for the heat equation is ill-posed. On the other hand, the ill-posedness of the periodic heat equation (1.1) is related to the fact that the set of eigenfunctions of the operator L does not form a basis in the Hilbert space $X = L^2_{\text{per}}([-\pi, \pi])$. Indeed, if the set of eigenfunctions forms a basis in X , then a dense operator L with a purely discrete spectrum of simple eigenvalues $\{\lambda_n\}_{n \in \mathbb{Z}}$ on $\text{Re } \lambda_n = 0$ is the infinitesimal generator of a strongly continuous semigroup.

To prove the last claim, we assume that the set of eigenfunctions $\{f_n\}_{n \in \mathbb{Z}}$ forms a basis in the Hilbert space X . Then, the solution of the inhomogeneous equation $(\lambda I - L)\phi = \psi$ with $\psi \in X$ can be solved by the series of eigenfunctions

$$\phi = \sum_{n \in \mathbb{Z}} \frac{(f_n^*, \psi)}{(f_n^*, f_n)} \frac{f_n}{\lambda - \lambda_n}, \quad \bar{\phi} = \sum_{n \in \mathbb{Z}} \frac{(f_n, \bar{\psi})}{(f_n^*, f_n)} \frac{\bar{f}_n^*}{\bar{\lambda} - \bar{\lambda}_n}. \tag{5.2}$$

Using the orthogonality of eigenfunctions f_n with respect to the adjoint eigenfunctions f_n^* and the normalization $(f_n^*, f_n) = 1$ for any $n \in \mathbb{Z}$, we obtain

$$\|\phi\|_X^2 = \sum_{n \in \mathbb{Z}} \frac{(f_n^*, \psi)(f_n, \bar{\psi})}{(\lambda - \lambda_n)(\bar{\lambda} - \bar{\lambda}_n)}, \quad \|\psi\|_X^2 = \sum_{n \in \mathbb{Z}} (f_n^*, \psi)(f_n, \bar{\psi}).$$

Since $\text{Re } \lambda_n = 0$ and $\text{Re } \lambda > 0$, then $\|\phi\|_X^2 \leq (\text{Re } \lambda)^{-2} \|\psi\|_X^2$, such that $\|(\lambda I - L)^{-1}\|_{X \rightarrow X} \leq 1/\text{Re } \lambda$ and the conditions of the Hille–Yosida Theorem are satisfied.

We conclude with the help of numerical computations that the Cauchy problem for the periodic heat equation (1.1) is ill-posed. Depending on the initial data h_0 , the blow-up of solutions may occur in finite or infinitesimal time, according to the conjecture in [3]. Although the series of eigenfunctions of operator L cannot be used to solve the Cauchy problem for the periodic heat equation, conditional convergence of the series of eigenfunctions can sometimes be achieved at least for finite times, as illustrated in [4]. Therefore, more detailed studies of applicability of the series of eigenfunctions and its dependence from the initial data h_0 are opened for further work.

Note. When the project was essentially complete, we became aware of a recent work [6], where similar results were obtained. In particular, the author of [6] proves that the spectral problem (1.3) has no continuous spectrum and, assisted with the numerical computations, illustrates that the eigenfunctions for isolated eigenvalues do not form a basis. The analysis of [6] is based on the difference eigenvalue problem (4.6), which makes it different from analysis in our work.

Acknowledgments

The authors thank E. Benilov for formulation of the problem, V. Ivrii and V. Strauss for useful discussions, and E.B. Davies for critical reading of our manuscript. M.C. is supported by the NSERC Graduate Fellowship. D.P. is supported by the Humboldt and EPSRC Research Fellowships. The numerical work was made possible by the facilities of the Shared Hierarchical Academic Research Computing Network (SHARCNET).

Appendix A. Spectrum of the linear operator L in weighted spaces

The operator L can be rewritten in the Sturm–Liouville symmetric form

$$L = -\epsilon \left| \cot\left(\frac{\theta}{2}\right) \right|^{1/\epsilon} L_0, \quad L_0 = \frac{d}{d\theta} \left(\left| \tan\left(\frac{\theta}{2}\right) \right|^{1/\epsilon} \sin \theta \frac{d}{d\theta} \right). \tag{A.1}$$

Let $r(\theta) = |\tan(\theta/2)|^{1/\epsilon}$ be the weight of the Sturm–Liouville spectral problem

$$-\epsilon L_0 f(\theta) = \lambda r(\theta) f(\theta), \tag{A.2}$$

acting on smooth functions $f(\theta)$ on $\theta \in [-\pi, \pi]$ in the weighted space $f \in L^2_r([-\pi, \pi])$.

Proposition A.1. *The operator L_0 admits a self-adjoint extension in $L^2_r([-\pi, \pi])$ for $0 < \epsilon < 1$, such that the spectrum of L_0 is purely discrete, consists of a set of simple real eigenvalues $\{\lambda_n\}_{n \in \mathbb{Z}}$ with $\lambda_0 = 0$, $\lambda_n = -\lambda_{-n}$, $\forall n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} \lambda_n = \infty$.*

Proof. By Lemma 6, the eigenfunction $f(\theta)$ of the spectral problem (A.2) for $0 < \epsilon < 1$ is in $L^2_r([-\pi, \pi])$ if and only if it is spanned by the fundamental solutions $f_1(\theta)$, $f_2^+(\theta)$ and $f_2^-(\theta)$, such that $f(\theta)$ is bounded at $\theta = 0$ and $\lim_{\theta \rightarrow \pm\pi} f(\theta) = 0$. Let $f(\theta)$ on $\pm\theta \in [0, \pi]$ be represented by

$$f(\theta) = \left[\pm \cot\left(\frac{\theta}{2}\right) \right]^{1/2\epsilon} g_{\pm}(x), \quad \cos \theta = x, \tag{A.3}$$

where $x \in [-1, 1]$. Then, the spectral problem (1.3) is rewritten in the form

$$-\epsilon \frac{d}{dx} \left[(1-x^2) \frac{dg_{\pm}}{dx} \right] + \frac{g_{\pm}(x)}{4\epsilon(1-x^2)} = \pm \frac{\lambda g_{\pm}(x)}{\sqrt{1-x^2}}. \tag{A.4}$$

If $f(\theta)$ belongs to $L^2_r([-\pi, \pi])$ for $0 < \epsilon < 1$, then $g_{\pm}(x)$ belong to $L^2_{\rho}([-1, 1])$ with the weight function $\rho(x) = \frac{1}{\sqrt{1-x^2}}$. Since the symmetric spectral problem (A.4) is self-adjoint in space $L^2_{\rho}([-1, 1])$, its eigenvalues λ are all real-valued. By Lemma 2 and the Sturm–Liouville theory [15], the eigenvalues $\{\lambda_n\}_{n \in \mathbb{Z}}$ are all simple and symmetric with $\lambda_0 = 0$ and $\lambda_n = -\lambda_{-n}$, $\forall n \in \mathbb{N}$, while the sequence $\{\lambda_n\}_{n \in \mathbb{Z}}$ is unbounded with $\lim_{n \rightarrow \infty} \lambda_n = \infty$. By the standard Green’s identity,

$$\lambda \|g_+\|_{L^2_{\rho}}^2 = \epsilon \int_{-1}^1 (1-x^2) |g'_+(x)|^2 dx + \int_{-1}^1 \frac{|g_+(x)|^2 dx}{4\epsilon(1-x^2)} > 0, \tag{A.5}$$

the eigenvalues λ_n of $g_+(x)$ are proved to be positive. If $\lambda_n > 0$ is an eigenvalue for $g_+(x)$, then $-\lambda_n < 0$ cannot be an eigenvalue for $g_-(x)$, such that $g_-(x) = 0$ on $x \in [-1, 1]$. Therefore, $f_n(\theta) = 0$ on $\theta \in [-\pi, 0]$ if $\lambda_n > 0$. The proof that the spectrum of L_0 in $L^2_r([-\pi, \pi])$ is purely discrete is done similarly to the proof of Lemma 5. \square

The operator L_0 admits a non-unique self-adjoint extension in $L^2_r([-\pi, \pi])$ for $\epsilon > 1$. Indeed, if $\epsilon > 1$, the eigenfunction $f(\theta)$ of the spectral problem (A.2) may exist in $L^2_r([-\pi, \pi])$ even if it is spanned by both fundamental solutions of Lemma 6 in the form

$$f(\theta) = Af_1(\theta) + Bf_2(\theta) = A^+ f_1^+(\theta) + B^+ f_2^+(\theta) = A^- f_1^-(\theta) + B^- f_2^-(\theta).$$

It follows from the Sturm–Liouville problem (A.2) that

$$\lambda \int_{-\pi}^{\pi} r(\theta) |f(\theta)|^2 d\theta = -\epsilon \left| \tan\left(\frac{\theta}{2}\right) \right|^{1/\epsilon} \sin(\theta) \bar{f}(\theta) f'(\theta) \Big|_{\theta=-\pi}^{\theta=\pi} + \epsilon \int_{-\pi}^{\pi} \left| \tan\left(\frac{\theta}{2}\right) \right|^{1/\epsilon} \sin(\theta) |f'(\theta)|^2 d\theta.$$

The second integral is finite for $\epsilon > 0$ only if $B = 0$. In this case, the first term is computed explicitly as follows

$$2^{1/\epsilon} (\bar{A}^+ B^+ + \bar{A}^- B^-). \tag{A.6}$$

This term is zero if $A^+ = A^- = 0$, which corresponds to the self-adjoint extension constructed in Proposition A.1. However, this choice is not unique, e.g. the alternative pairing $B^+ = B^- = 0$ can also be applied.

Appendix B. Resonant poles of the Schrödinger operators

Eigenvalues of the operator L coincide for $|\epsilon| > \frac{1}{2}$ with resonant poles of the Schrödinger operators. To show this, we use the transformation (3.5) on the intervals $\pm\theta \in [0, \pi]$ and rewrite the spectral problem (1.3) as the uncoupled spectral problems (3.6) for $f(\theta) \equiv f_{\pm}(t)$ on $t \in \mathbb{R}$. Let $f(\theta)$ satisfy the normalization condition $f(\pi) = f(-\pi) = 1$ and $\lambda \notin \mathbb{R}$. Then, eigenfunctions $f_{\pm}(t)$ of the uncoupled problems (3.6) satisfy the boundary conditions

$$\lim_{t \rightarrow -\infty} f_{\pm}(t) = 1, \quad \lim_{t \rightarrow \infty} f_{\pm}(t) = a^{\pm}, \tag{B.1}$$

where a^{\pm} are uniquely defined. The function $f(\theta)$ on $\theta \in [-\pi, \pi]$ constructed from $f_{\pm}(t)$ on $t \in \mathbb{R}$ is continuous at $\theta = 0$ if $a^+ = a^-$.

Let $\epsilon > \frac{1}{2}$ and define $f_{\pm}(t) = e^{\frac{t}{2\epsilon}} g_{\pm}(t)$. The eigenfunctions $g_{\pm}(t)$ satisfy the linear Schrödinger equations

$$\left(\frac{1}{4\epsilon} - \epsilon \partial_t^2\right) g_{\pm} = \pm \lambda \operatorname{sech} t g_{\pm}, \tag{B.2}$$

but the boundary conditions (B.1) show that $g_{\pm} \notin L^2(\mathbb{R})$ and λ is not an eigenvalue. In fact, the eigenfunctions $g_{\pm}(t)$ belong to the exponentially weighted L^2 -space, such that λ is a resonance pole of the Schrödinger operators. Let us decompose the eigenfunctions by $g_{\pm}(t) = e^{-\frac{t}{2\epsilon}} + h_{\pm}(t)$. By the theory of exponential asymptotics of solutions of the Schrödinger problems (B.2), it follows that $h_{\pm} \in L^2(\mathbb{R})$ if $\epsilon > \frac{1}{2}$. Let $h_0(t) = e^{-\frac{t}{2\epsilon}} \operatorname{sech} t \in L^2(\mathbb{R})$ and define the linear inhomogeneous problems for $h_{\pm}(t)$,

$$S_{\epsilon}^{\pm}(\lambda)h_{\pm} = \pm \lambda h_0(t), \quad S_{\epsilon}^{\pm}(\lambda) = \frac{1}{4\epsilon} - \epsilon \partial_t^2 \mp \lambda \operatorname{sech} t. \tag{B.3}$$

The operator $S_{\epsilon}^{\pm}(\lambda)$ maps $H^2(\mathbb{R})$ to $L^2(\mathbb{R})$ and, if $\lambda \notin \mathbb{R}$, the kernel of S_{ϵ}^{\pm} is empty. The boundary condition $a^+ = a^-$ is then equivalent to the zeros of the function

$$H_{\epsilon}(\lambda) = \lambda \lim_{t \rightarrow \infty} e^{\frac{t}{2\epsilon}} [(S_{\epsilon}^+)^{-1}(\lambda) + (S_{\epsilon}^-)^{-1}(-\lambda)] h_0(t). \tag{B.4}$$

The function $H_{\epsilon}(\lambda)$ coincides (up to a multiplicative constant) with the analytic function $F_{\epsilon}(\lambda)$ introduced in Corollary 1 for $|\epsilon| < 2$. Therefore, $H_{\epsilon}(\lambda)$ represents a continuation of $F_{\epsilon}(\lambda)$ from the domain $|\epsilon| < 2$ to the domain $|\epsilon| > \frac{1}{2}$. The function $H_{\epsilon}(\lambda)$ is analytic in $\lambda \in \mathbb{C}$ and its roots give isolated eigenvalues of the spectral problem (1.3) with the account of their multiplicity.

The function $H_{\epsilon}(\lambda)$ can be simplified for $\lambda \in i\mathbb{R}$. Let $\lambda = i\omega \in i\mathbb{R}$ and represent $h_{\pm} = F(t) \pm iG(t)$, where

$$L_{\epsilon}F = -\omega \operatorname{sech} t G, \quad L_{\epsilon}G = \omega \operatorname{sech} t F + \omega h_0(t), \quad L_{\epsilon} = \frac{1}{4\epsilon} - \epsilon \partial_t^2. \tag{B.5}$$

Therefore, we can define a real-valued function $\tilde{H}_{\epsilon}(\omega) = H_{\epsilon}(i\omega)$ on $\omega \in \mathbb{R}$ given by

$$\tilde{H}_{\epsilon}(\omega) = \omega \lim_{t \rightarrow \infty} e^{\frac{t}{2\epsilon}} (L_{\epsilon} + \omega^2 \operatorname{sech} t L_{\epsilon}^{-1} \operatorname{sech} t)^{-1} h_0(t). \tag{B.6}$$

By multiplying the linear inhomogeneous equation

$$(L_{\epsilon} + \omega^2 \operatorname{sech} t L_{\epsilon}^{-1} \operatorname{sech} t)G = \omega h_0(t),$$

by $e^{\frac{t}{2\epsilon}}$ and integrating on $t \in \mathbb{R}$ by parts, the function $\tilde{H}_{\epsilon}(\omega)$ can be represented in the integral form

$$\tilde{H}_{\epsilon}(\omega) = \omega \int_{-\infty}^{\infty} \operatorname{sech} t dt - \omega^2 \int_{-\infty}^{\infty} e^{\frac{t}{2\epsilon}} \operatorname{sech} t L_{\epsilon}^{-1} \operatorname{sech} t G(t) dt.$$

Let $\tilde{h}_0(t) = \operatorname{sech} t L_{\epsilon}^{-1} \operatorname{sech} t e^{\frac{t}{2\epsilon}}$. Then,

$$\tilde{H}_{\epsilon}(\omega) = \pi\omega - \omega^2 \int_{-\infty}^{\infty} \tilde{h}_0(t)G(t) dt.$$

Zeros of the real analytic function $\tilde{H}_{\epsilon}(\omega)$ are equivalent to purely imaginary eigenvalues of the spectral problem (1.3) for $|\epsilon| > \frac{1}{2}$.

References

[1] N.I. Akhiezer, I.M. Glazman, Theory of Linear Operators in Hilbert Space, Frederic Ungar Publishing Co., New York, 1963.
 [2] F.V. Atkinson, Discrete and Continuous Boundary Problems, Academic Press, London, 1964.
 [3] E.S. Benilov, S.B.G. O'Brien, I.A. Sazonov, A new type of instability: Explosive disturbances in a liquid film inside a rotating horizontal cylinder, J. Fluid Mech. 497 (2003) 201–224.
 [4] E.S. Benilov, Explosive instability in a linear system with neutrally stable eigenmodes. Part 2. Multi-dimensional disturbances, J. Fluid Mech. 501 (2004) 105–124.

- [5] E.B. Davies, *Spectral Theory and Differential Operators*, Cambridge Univ. Press, 1995.
- [6] E.B. Davies, An indefinite convection–diffusion operator, *LMS J. Comput. Math.* 10 (2007) 288–306.
- [7] E.B. Davies, M. Plum, Spectral pollution, *IMA J. Numer. Anal.* 24 (2004) 417–438.
- [8] N. Dunford, J.T. Schwartz, *Linear Operators. Part II: Spectral Theory*, John Wiley & Sons, New York, 1963.
- [9] V. Hernandez, J.E. Roman, V. Vidal, A scalable and flexible toolkit for the solution of eigenvalue problems, *ACM Trans. Math. Software* 31 (2005).
- [10] I.M. Glazman, *Direct Methods of Qualitative Spectral Analysis of Singular Differential Operators*, Israel Program for Scientific Translations Ltd., 1965.
- [11] I. Gohberg, S. Goldberg, M.A. Kaashoek, *Classes of Linear Operators*, vol. 1, Birkhäuser Verlag, Basel, 1990.
- [12] I. Gohberg, M.G. Krein, *Introduction to the Theory of Linear Non-Selfadjoint Operators*, Amer. Math. Soc. Transl., vol. 18, Amer. Math. Soc., Providence, RI, 1969.
- [13] M. Lavrentjev, L. Saveljev, *Linear Operators and Ill-Posed Problems*, Consultants Bureau, New York, 1995.
- [14] J.T. Marti, *Introduction to the Theory of Bases*, Springer-Verlag, New York, 1969.
- [15] P.M. Morse, H. Feshbach, *Methods of Theoretical Physics*, vol. 1, McGraw–Hill Book Co., New York, 1953.
- [16] Y. Saad, *Numerical Methods for Large Eigenvalue Problems*, Manchester Univ. Press, Manchester, 1992.
- [17] L.N. Trefethen, M. Embree, *Spectra and Pseudospectra. The Behavior of Nonnormal Matrices and Operators*, Princeton Univ. Press, Princeton, NJ, 2005.
- [18] K. Yosida, *Functional Analysis*, Springer-Verlag, New York, 1968.