

# NORMAL FORM FOR TRANSVERSE INSTABILITY OF THE LINE SOLITON WITH A NEARLY CRITICAL SPEED OF PROPAGATION

DMITRY PELINOVSKY

ABSTRACT. There exists a critical speed of propagation of the line solitons in the Zakharov–Kuznetsov (ZK) equation such that small transversely periodic perturbations are unstable for line solitons with larger-than-critical speeds and orbitally stable for those with smaller-than-critical speeds. The normal form for transverse instability of the line soliton with a nearly critical speed of propagation is derived by means of symplectic projections and near-identity transformations. Justification of this normal form is provided with the energy method. The normal form predicts a transformation of the unstable line solitons with larger-than-critical speeds to the orbitally stable transversely modulated solitary waves.

## 1. INTRODUCTION

Starting with the pioneer work of Kadomtsev & Petviashvili [6] and Zakharov & Rubenchik [31], it is well known that the line solitons are spectrally unstable with respect to the long transverse perturbations in many nonlinear evolution equations such as the Kadomtsev–Petviashvili (KP) and nonlinear Schrödinger (NLS) equations (see review in [9]). The spectral instability persists to the short transverse perturbations of any period in the hyperbolic version of the two-dimensional NLS equation [4, 22], whereas it disappears for short transverse perturbations in many other equations such as the elliptic version of the two-dimensional NLS equation and the KP-I equation [23, 24]. Alternatively, for a fixed period of the transverse perturbation, the transverse instability occurs for the line solitons with larger-than-critical speeds of propagation and disappears for those with smaller-than-critical speeds.

In the prototypical case of the KP-I equation, it was shown in [25] that if the line solitons are spectrally stable with respect to the periodic perturbations, they remain nonlinearly stable, whereas if they are spectrally unstable, they remain nonlinearly unstable. The spectral stability analysis is inconclusive for the line soliton with the critical speed of propagation since the linearized operator has an additional zero eigenvalue beyond the one induced by the translational symmetry of the KP-I equation. The presence of the additional zero eigenvalue implies a bifurcation of the new travelling solutions which are spatially localized along the longitudinal direction and are periodic along the transverse perturbations. Such travelling solitary waves with periodic transverse modulations were discovered for the KP-I equation by Zaitsev [28].

Analytical solutions for the unstable eigenmode were derived for the KP-I equation by Zakharov [29]. Exact solutions for the nonlinear evolution of the unstable line solitons were obtained and analyzed in [15, 20]. If the transverse perturbation is proportional to

---

*Date:* May 31, 2017.

The results of this work were obtained with the financial support from the state task of Russian Federation in the sphere of scientific activity (Task No. 5.5176.2017/BCh).

a single unstable eigenmode, it results in the monotonic transformation of the unstable line soliton with a larger-than-critical speed of propagation to the travelling solitary wave with the periodic transverse modulation of the same period and an ejection of a stable line soliton with a smaller-than-critical speed. In the case of multi-mode perturbations, several modulated travelling waves and the residual line soliton are generated in the dynamics of an unstable line soliton, according to the exact solutions of the KP-I equation [20].

Transformation of the unstable line solitons with a nearly critical speed of propagation was studied in the framework of the KP-I equation with an asymptotic multi-scale expansion method [5]. An integrable Eckhaus equation was derived from the integrable KP-I equation. This Eckhaus equation correctly represents the monotonic transition of the unstable line soliton to the transversely modulated solitary wave and a “radiation” of a stable line soliton of a smaller speed of propagation, in comparison with the exact solutions to the KP-I equation (see Section 4.7 in [9]). Similar asymptotic reductions were reported in [21] for the line dark solitons of the defocusing elliptic NLS equation.

The present work is devoted to the justification of the asymptotic model describing the nonlinear dynamics of the transverse perturbations on the line soliton with a nearly critical speed of propagation. Stability of these line solitons and transversely modulated solitary waves are derived as a by-product of this asymptotic model. Unfortunately, the analytical setup does not apply to the KP-I equation, partly, because the continuous spectrum of the linearized operators does not move to the left-half plane in exponentially weighted spaces. For a better model, where this difficulty does not arise, we consider the Zakharov–Kuznetsov (ZK) equation,

$$(1.1) \quad u_t + 12uu_x + u_{xxx} + u_{xyy} = 0,$$

which features a similar phenomenon. The ZK equation is an anisotropic generalization of the Korteweg–De Vries (KdV) equation in two spatial dimensions [30]. Justification of the ZK equation in the context of the Euler–Poisson equations for magnetized plasmas was recently reported in [11]. Transverse stability of line solitons is very similar between the ZK and KP-I equations [23, 24], but the ZK equation can be analyzed successfully by using exponentially weighted Sobolev spaces, similar to the analysis of the KdV equation [16, 17]. Instability of line solitons in the ZK equation is known for quite some time in physics literature (see, e.g., [1]).

We shall derive and justify the first-order differential equation

$$(1.2) \quad \frac{db}{dt} = \lambda'(c_*)(c_+ - c_*)b + \gamma|b|^2b, \quad t > 0,$$

where  $\lambda'(c_*) > 0$ ,  $\gamma < 0$  are real-valued numerical coefficients,  $c_*$  is the critical speed of the line soliton,  $c_+ \in \mathbb{R}$  depends on the initial conditions, and  $b(t) : \mathbb{R}_+ \rightarrow \mathbb{C}$  is an amplitude of transverse perturbation. The differential equation (1.2) describes the nonlinear dynamics of a small transverse perturbation of a fixed period to the line soliton with a nearly critical speed  $c_*$  and is referred to as “*normal form for transverse instability of the line soliton with a nearly critical speed of propagation*”.

If  $c_+ < c_*$ , the normal form equation (1.2) describes a monotonic exponential decay  $b(t) \rightarrow 0$  as  $t \rightarrow +\infty$  and suggests that the line solitons with the smaller-than-critical speeds are asymptotically stable with respect to small transverse perturbations of a fixed period [27]. If  $c_+ \geq c_*$ , the normal form equation (1.2) describes a monotonic algebraic

decay  $b(t) \rightarrow b_*$  as  $t \rightarrow \infty$ , where

$$(1.3) \quad |b_*| = \sqrt{\frac{\lambda'(c_*)(c_+ - c_*)}{|\gamma|}}, \quad c_+ \geq c_*.$$

This solution suggests a transition from the unstable line solitons with the larger-than-critical speeds to stable transversely modulated solitary waves, which are also asymptotically stable with respect to small transverse perturbations of the same fixed period [27].

Bifurcations and stability of the line solitons under the transverse perturbations of a critical period were addressed recently by Yamazaki for the elliptic version of the NLS equation in [26] and for the ZK equation (1.1) in [27].

In [26], the bifurcation problem is analyzed with the Lyapunov–Schmidt reduction method and the nonlinear orbital stability is deduced from the energy method based on convexity of the action functional for the NLS equation, which is the same as for the ZK equation (1.1). Similarly to our work here, it is reported in [26] that the transversely modulated solitary waves are orbitally stable for the case of quadratic nonlinearities. Dynamics near such waves was not studied in [26].

In [27], asymptotic stability of the line solitons with the smaller-than-critical speeds and the transversely modulated solitary waves with the larger-than-critical speeds was shown for the ZK equation (1.1) with a Liouville-type theorem and virial type estimates.

Asymptotic stability of two-dimensional solitary waves was considered in the  $L^2$ -subcritical ZK equations [3], which includes the ZK equation (1.1).

The remainder of this paper is organized as follows. Section 2 contains results on transverse instability of line solitons and bifurcations of transversely modulated solitary waves in the ZK equation (1.1). Formal derivation and justification of the normal form (1.2) is given in Section 3.

**Acknowledgement.** This paper was mostly written in 2014–15 after discussions with F. Rousset (Paris–Sud) and N. Tzvetkov (Cergy–Pontoise) before papers [26] and [27] were first posted on arXiv and then published. The author thanks the collaborators for useful discussions and valuable comments.

## 2. TRANSVERSE INSTABILITY OF LINE SOLITONS FOR THE ZK EQUATION (1.1)

We are concerned here with the transverse instability of the line solitons under periodic transverse perturbations in the ZK equation (1.1). First, we review relevant properties of the line solitons of the KdV equation. Next, we obtain spectral transverse stability results for the line solitons with a nearly critical speed of propagation. Further, we study bifurcations of the transversely modulated solitary waves. Finally, we present the main result on the normal form for transverse instability of the line solitons with a nearly critical speed of propagation.

**2.1. Properties of line solitons of the KdV equation.** Line solitons are expressed analytically as the two-parameter family

$$(2.1) \quad u_c(\xi) = c \operatorname{sech}^2(\sqrt{c}\xi), \quad \xi = x - 4ct - x_0,$$

where  $c > 0$  is the speed parameter and  $x_0 \in \mathbb{R}$  is the translation parameter. The line soliton (2.1) for a fixed  $c > 0$  is a critical point of the action functional

$$(2.2) \quad \Lambda_c(u) := \frac{1}{2} \int_{\mathbb{R}} [(\partial_\xi u)^2 - 4u^3 + 4cu^2] d\xi,$$

which is a linear combination of the energy and momentum of the KdV equation. The second variation of the action functional (2.2) at the line soliton (2.1) is defined by the Hessian operator  $L_c : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ , the differential expression of which is given by

$$(2.3) \quad L_c := -\partial_\xi^2 + 4c - 12c \operatorname{sech}^2(\sqrt{c}\xi).$$

The Schrödinger operator  $L_c : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is known [10] to have the essential spectrum located on  $[4c, \infty)$  and three simple isolated eigenvalues  $\lambda_1 < \lambda_2 < \lambda_3 < 4c$ . More precisely, the eigenvalues and the corresponding non-normalized eigenfunctions of  $L_c$  are given by

$$(2.4) \quad \lambda_1 = -5c, \quad \varphi_1 = \operatorname{sech}^3(\sqrt{c}\xi),$$

$$(2.5) \quad \lambda_2 = 0, \quad \varphi_2 = \operatorname{sech}^2(\sqrt{c}\xi) \tanh(\sqrt{c}\xi),$$

$$(2.6) \quad \lambda_3 = 3c, \quad \varphi_3 = 4\operatorname{sech}(\sqrt{c}\xi) - 5\operatorname{sech}^3(\sqrt{c}\xi).$$

The first two eigenvalues and eigenfunctions are particularly important in the spectral transverse stability analysis of the line solitons (2.1).

**2.2. Spectral transverse stability analysis of line solitons.** Let us consider the  $2\pi$ -periodic transverse perturbation to the line solitons (2.1). Therefore, we fix the length of the transverse periodic perturbation and vary the speed parameter  $c$ . Substituting  $u(x, t) = u_c(\xi) + U(\xi)e^{\lambda t + iky}$  with  $k \in \mathbb{Z}$  into the ZK equation (1.1) and dropping the quadratic terms in  $U$  yields the spectral problem

$$(2.7) \quad \partial_\xi(L_c + k^2)U = \lambda U, \quad k \in \mathbb{Z},$$

where  $k$  is the wave number of the  $2\pi$ -periodic transverse perturbation and  $\partial_\xi L_c : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is the linearized operator for the KdV equation.

For  $k = 0$ , the spectral problem (2.7) coincides with the one for the KdV equation. It is known from the work of Pego & Weinstein [16] that the spectrum of  $\partial_\xi L_c : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  consists of a double zero eigenvalue and a continuous spectrum on  $i\mathbb{R}$ . The double zero eigenvalue is associated with the following Jordan block of the operator  $\partial_\xi L$ :

$$(2.8) \quad \partial_\xi L_c \partial_\xi u_c = 0, \quad \partial_\xi L_c \partial_c u_c = -4\partial_\xi u_c,$$

where the derivatives of  $u_c$  in  $\xi$  and  $c$  are exponentially decaying functions of  $\xi$ . The following lemma characterizes the spectral problem (2.7) for any  $k \in \mathbb{N}$ .

**Lemma 2.1.** *For any  $k \in \mathbb{N}$ , the spectral problem (2.7) has a pair of real eigenvalues  $\pm\lambda_k(c)$  if  $c > c_k := \frac{k^2}{5}$ . No eigenvalues with  $\operatorname{Re}(\lambda) \neq 0$  exist if  $c \in (0, c_k)$ .*

**Proof.** For any  $k \in \mathbb{N}$ , the self-adjoint Schrödinger operator  $L_c + k^2 : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is strictly positive for  $c \in (0, c_k)$  and admits a simple negative eigenvalue for  $c > c_k$ , where  $c_k := \frac{k^2}{5}$ . It follows from the main theorem in [7, 18] that the spectrum of  $\partial_\xi(L_c + k^2) : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  has exactly one pair of real eigenvalues  $\pm\lambda_k(c)$  if  $c > c_k$  and no eigenvalues with  $\operatorname{Re}(\lambda) \neq 0$  if  $c \in (0, c_k)$ .  $\blacksquare$

**Remark 2.1.** *By using the energy method in [25], one can easily prove nonlinear orbital stability of the line solitons with  $c \in (0, c_*)$  and nonlinear instability of the line solitons with  $c > c_*$ , where*

$$c_* := \min_{k \in \mathbb{N}} c_k \equiv \frac{1}{5}.$$

The following result gives a precise characterization of the unstable eigenvalue bifurcating at  $c = c_*$  in the spectral problem (2.7) with  $k = 1$ . To do so, we introduce the exponentially weighted space

$$(2.9) \quad H_\mu^s(\mathbb{R}) = \{u \in H_{\text{loc}}^s(\mathbb{R}) : e^{\mu\xi}u \in H^s(\mathbb{R})\}, \quad s \geq 0, \quad \mu > 0.$$

This weighted space is used to push the continuous spectrum of the operator  $\partial_\xi L_c : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  to the left-half plane for  $\mu > 0$  sufficiently small [16, 17]. In the generalized KdV equation with a triple zero eigenvalue, the exponentially weighted space is used to construct the Jordan block for the triple zero eigenvalue in [2, 13].

**Lemma 2.2.** *There is  $\mu_0 > 0$  such that for every  $\mu \in (0, \mu_0)$ , the spectral problem (2.7) with  $k = 1$  and  $c = c_*$  considered in  $L_\mu^2(\mathbb{R})$  admits a simple zero eigenvalue with the eigenfunction  $\psi_* \in H_\mu^3(\mathbb{R})$  and the adjoint eigenfunction  $\eta_* \in H_{-\mu}^3(\mathbb{R})$ , where*

$$(2.10) \quad \psi_*(\xi) = \text{sech}^3(\sqrt{c_*}\xi), \quad \eta_*(\xi) = \int_{-\infty}^{\xi} \text{sech}^3(\sqrt{c_*}\xi')d\xi',$$

hence,

$$(2.11) \quad \langle \eta_*, \psi_* \rangle_{L^2} = \frac{1}{2} \left( \int_{\mathbb{R}} \text{sech}^3(\sqrt{c_*}\xi)d\xi \right)^2 = \frac{\pi^2}{8c_*}.$$

Moreover, for a given  $\mu \in (0, \mu_0)$ , there exist an interval  $(c_-, c_+)$  with  $c_- < c_* < c_+$  such that the spectral problem (2.7) with  $k = 1$  and  $c \in (c_-, c_+)$  considered in  $L_\mu^2(\mathbb{R})$  admits a small eigenvalue  $\lambda(c)$ , where the mapping  $c \mapsto \lambda$  is smooth and is given by

$$(2.12) \quad \lambda(c) = \lambda'(c_*)(c - c_*) + \mathcal{O}((c - c_*)^2) \quad \text{as } c \rightarrow c_*,$$

with

$$(2.13) \quad \lambda'(c_*) = \frac{128}{3\pi^2} \sqrt{c_*}.$$

**Proof.** Existence of the zero eigenvalue of  $\partial_\xi(L_{c_*} + 1) : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  follows from existence of the negative eigenvalue  $-1$  of  $L_{c_*}$  in (2.4). The eigenfunction  $\psi_* = \varphi_1$  at  $c = c_*$  is exponentially decaying in  $\xi$ , therefore, there is a positive  $\mu_0$  such that  $e^{\mu\xi}\psi_*$  is exponentially decaying in  $\xi$  for every  $\mu \in (0, \mu_0)$ . Thus, the operator  $\partial_\xi(L_{c_*} + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  has a simple zero eigenvalue.

If  $H_\mu^3(\mathbb{R})$  is the domain of  $\partial_\xi(L_{c_*} + 1)$ , then  $H_{-\mu}^3(\mathbb{R})$  is the domain of the adjoint operator  $-(L_{c_*} + 1)\partial_\xi$ . For  $\mu > 0$ , the eigenfunction  $\eta_*$  of the kernel of  $-(L_{c_*} + 1)\partial_\xi : H_{-\mu}^3(\mathbb{R}) \rightarrow L_{-\mu}^2(\mathbb{R})$  must decay faster than  $e^{-\mu\xi}$  grows as  $\xi \rightarrow -\infty$ . Since  $\partial_\xi\eta_*$  is proportional to  $\psi_*$  and  $\eta_*(\xi) \rightarrow 0$  as  $\xi \rightarrow -\infty$ , we set

$$\eta_*(\xi) := \int_{-\infty}^{\xi} \psi_*(\xi')d\xi'$$

and obtain (2.10). The nonzero inner product in (2.11) ensures that the Jordan block for the zero eigenvalue of  $\partial_\xi(L_{c_*} + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  with  $\mu \in (0, \mu_0)$  is simple.

It remains to prove the last assertion of the lemma with the expansion (2.12). Since the simple zero eigenvalue of  $\partial_\xi(L_c + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  at  $c = c_*$  is isolated from the continuous spectrum of this operator for any fixed  $\mu \in (0, \mu_0)$ , one can use analytic perturbation theory [8]. In particular, the operator  $L_c : H^2(\mathbb{R}) \mapsto L^2(\mathbb{R})$  is analytic at  $c = c_*$  and admits the following expansion

$$(2.14) \quad L_c = L_{c_*} + L'_{c_*}(c - c_*) + \tilde{L}_c(c - c_*)^2,$$

where

$$(2.15) \quad L'_{c_*} = 4 - 12\operatorname{sech}^2(\sqrt{c_*}\xi) + 12\sqrt{c_*}\xi \operatorname{sech}^2(\sqrt{c_*}\xi) \tanh(\sqrt{c_*}\xi)$$

and  $\tilde{L}_c$  is an exponentially decaying and bounded potential as  $c \rightarrow c_*$ . By using perturbation expansions,

$$(2.16) \quad \lambda(c) = \lambda_1(c - c_*) + \mathcal{O}((c - c_*)^2), \quad \psi(c) = \psi_* + \psi_1(c - c_*) + \mathcal{O}_{H_\mu^3}((c - c_*)^2),$$

we obtain the linear inhomogeneous equation at the order of  $\mathcal{O}(c - c_*)$ :

$$(2.17) \quad \partial_\xi(L_{c_*} + 1)\psi_1 + \partial_\xi L'_{c_*}\psi_* = \lambda_1\psi_*.$$

This equation is considered in  $L_\mu^2(\mathbb{R})$  for a fixed  $\mu \in (0, \mu_0)$ . Projecting (2.17) to  $\eta_*$ , the eigenfunction in the kernel of the adjoint operator  $-(L_{c_*} + 1)\partial_\xi$  in  $L_{-\mu}^2(\mathbb{R})$ , we obtain

$$(2.18) \quad \lambda_1 \langle \eta_*, \psi_* \rangle_{L^2} = \langle \eta_*, \partial_\xi L'_{c_*}\psi_* \rangle_{L^2} = -\langle \psi_*, L'_{c_*}\psi_* \rangle_{L^2}.$$

From (2.11), (2.15), and (2.18), we obtain

$$\langle \psi_*, L'_{c_*}\psi_* \rangle_{L^2} = -\frac{16}{3\sqrt{c_*}} \quad \text{hence} \quad \lambda_1 = \frac{128}{3\pi^2}\sqrt{c_*},$$

which agrees with (2.13) because  $\lambda_1 = \lambda'(c_*)$ . Justification of the asymptotic expansion (2.16) is developed with the analytic perturbation theory (Theorem 1.7 in [8, Chapter VII]).  $\blacksquare$

**Remark 2.2.** *Since the continuous spectrum of  $\partial_\xi(L_c + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  is located in the left-half plane for  $\mu > 0$  [16], whereas the eigenvalue  $\lambda(c)$  is positive for  $c > c_*$ , we can send  $\mu \rightarrow 0^+$  without affecting the eigenvalue  $\lambda(c) > 0$ . Therefore, the eigenvalue  $\lambda(c)$  persists for the operator  $\partial_\xi(L_c + 1) : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ . By symmetry  $\xi \mapsto -\xi$  and  $\lambda \mapsto -\lambda$ , the operator  $\partial_\xi(L_c + 1) : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  also admits the eigenvalue  $-\lambda(c) < 0$  for the same case  $c > c_*$ . Thus, the spectral problem (2.7) with  $k = 1$  has a pair of real eigenvalues  $\pm\lambda(c)$  for  $c > c_*$ , in agreement with Lemma 2.1.*

**Remark 2.3.** *Since the eigenvalue  $\lambda(c)$  in  $L_\mu^2(\mathbb{R})$  is negative for  $c < c_*$ , we cannot send  $\mu \rightarrow 0^+$  without affecting the eigenvalue  $\lambda(c) < 0$  by the deformation of the continuous spectrum of  $\partial_\xi(L_c + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$ . As is well-known [16], the eigenvalue  $\lambda(c) < 0$  of the operator  $\partial_\xi(L_c + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  becomes a resonant pole of the operator  $\partial_\xi(L_c + 1) : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  for  $c < c_*$ .*

**Remark 2.4.** *Perturbation expansions (2.14) and (2.16) can also be used to characterize the shift of the zero eigenvalue of the operator  $L_c + 1$  in  $L^2(\mathbb{R})$  for  $c \neq c_*$ . In this case,  $\psi_1$  satisfies*

$$(L_{c_*} + 1)\psi_1 + L'_{c_*}\psi_* = \lambda_1\psi_1,$$

from which we obtain

$$\lambda_1\|\psi_*\|_{L^2}^2 = \langle \psi_*, L'_{c_*}\psi_* \rangle_{L^2}.$$

Since  $\langle \psi_*, L'_{c_*}\psi_* \rangle_{L^2} < 0$ , we have  $\lambda_1 < 0$ . Therefore, the zero eigenvalue of the operator  $L_{c_*} + 1$  in  $L^2(\mathbb{R})$  becomes a negative eigenvalue of this operator for  $c > c_*$ . This shift induces the spectral instability of the line soliton, in agreement with Lemma 2.1.

**2.3. Transversely modulated solitary waves.** The instability bifurcation of the line solitons in Lemma 2.2 is related to the bifurcation of a new family of travelling solitary waves with the periodic transverse modulation. Such transversely modulated solitary waves satisfy the nonlinear elliptic problem

$$(2.19) \quad -\frac{\partial^2 u}{\partial \xi^2} - \frac{\partial^2 u}{\partial y^2} + 4cu - 6u^2 = 0, \quad (x, y) \in \mathbb{R} \times \mathbb{T},$$

where  $\mathbb{T}$  is a  $2\pi$ -periodic torus. In order to eliminate translational symmetries in  $\xi$  and  $y$ , we define the space of even functions both in  $\xi$  and  $y$ :

$$(2.20) \quad H_{\text{even}}^s = \{u \in H^s(\mathbb{R} \times \mathbb{T}) : u(-\xi, y) = u(\xi, y) = u(\xi, -y)\}, \quad s \geq 0.$$

The following lemma describes bifurcation of the transversely modulated solitary waves from the line soliton with a nearly critical speed of propagation.

**Lemma 2.3.** *There exists  $c_+ > c_*$  such that for every  $c \in (c_*, c_+)$ , the nonlinear elliptic problem (2.19) has a nontrivial solution  $u_b$  in  $H_{\text{even}}^2$  in addition to the line soliton (2.1). The solution  $u_b$  is expressed by the expansion*

$$(2.21) \quad u_b(\xi, y) = u_{c_*}(\xi) + 2b \cos(y)\psi_*(\xi) + \tilde{u}_b(\xi, y),$$

where  $b \in \mathbb{R}$  is a nonzero root of the algebraic equation

$$(2.22) \quad \alpha(c - c_*)b + \beta b^3 = 0$$

and  $\tilde{u}_b \in H_{\text{even}}^2$  satisfies the bound  $\|\tilde{u}_b\|_{H^2} \leq Ab^2$  for a positive constant  $A$  independently of  $b$  and  $c$ . Here

$$\alpha = -\langle \psi_*, L'_{c_*}\psi_* \rangle_{L^2} = \frac{16}{3\sqrt{c_*}} > 0$$

and  $\beta < 0$  is a numerical coefficient given by (2.38) below.

**Proof.** The algorithm follows the method of Lyapunov–Schmidt reduction and it is relatively well known, see, e.g., [19]. We write the decomposition (2.21) in  $H_{\text{even}}^2$  equipped with the orthogonality condition  $\langle v_1, \tilde{u}_b \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = 0$ , where  $v_1(\xi, y) := \cos(y)\psi_*(\xi)$  is the eigenfunction of the kernel of  $L_{c_*} - \partial_y^2 : H_{\text{even}}^2 \rightarrow L_{\text{even}}^2$ . For every  $c = c_* + \delta$  with  $\delta \in \mathbb{R}$  being sufficiently small, the correction term  $\tilde{u}_b \in H_{\text{even}}^2$  and the parameter  $b \in \mathbb{R}$  are defined by the projection equations of the Lyapunov–Schmidt reduction method:

$$(2.23) \quad (L_{c_*} - \partial_y^2 + 4\delta)\tilde{u}_b = -4\delta u_{c_*} - 8\delta b \cos(y)\psi_* + 6(2b \cos(y)\psi_* + \tilde{u}_b)^2$$

and

$$(2.24) \quad -4\delta b \|\psi_*\|_{L^2(\mathbb{R})}^2 + \frac{3}{\pi} \langle \cos(y)\psi_*, (2b \cos(y)\psi_* + \tilde{u}_b)^2 \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = 0.$$

Due to the bifurcation equation (2.24) and the symmetry constraints in (2.20), the linear operator in the left-hand side of equation (2.23) can be inverted with a bounded inverse in  $L^2_{\text{even}}$  for every small  $\delta \in \mathbb{R}$ . As a result, for every small  $b \in \mathbb{R}$  and  $\delta \in \mathbb{R}$ , the fixed-point argument can be applied to solve equation (2.23) and to obtain a unique  $\tilde{u}_b \in H^2_{\text{even}}$  satisfying the bound

$$(2.25) \quad \|\tilde{u}_b\|_{H^2} \leq A(|\delta| + b^2),$$

where the positive constant  $A$  is independent of  $\delta$  and  $b$ . However, the solution  $\tilde{u}_b \in H^2_{\text{even}}$  with the bound (2.25) is insufficient for the derivation of the algebraic equation (2.22) from the bifurcation equation (2.24) [19].

In order to obtain the algebraic equation (2.22), we perform a near-identity transformation

$$(2.26) \quad \tilde{u}_b(\xi, y) = 2b^2 \cos(2y)w_2(\xi) + b^2w_0(\xi) + \delta \partial_c u_{c_*}(\xi) + w(\xi, y),$$

where

$$(2.27) \quad \partial_c u_{c_*}(\xi) = \text{sech}^2(\sqrt{c_*}\xi) - \sqrt{c_*}\xi \tanh(\sqrt{c_*}\xi) \text{sech}^2(\sqrt{c_*}\xi),$$

the correction terms  $w_0$  and  $w_2$  are found from the linear inhomogeneous equations:

$$(2.28) \quad L_{c_*} w_0 = 12\psi_*^2$$

and

$$(2.29) \quad (L_{c_*} + 4)w_2 = 6\psi_*^2,$$

whereas  $w$  satisfies the transformed equation

$$(2.30) \quad \begin{aligned} (L_{c_*} - \partial_y^2 + 4\delta)w &= -4\delta^2 \partial_c u_{c_*} - 8\delta b^2 \cos(2y)w_2 - 4\delta b^2 w_0 \\ &\quad - 8\delta b \cos(y)\psi_* + 24b \cos(y)\psi_* \tilde{u}_b + 6\tilde{u}_b^2, \end{aligned}$$

where  $\tilde{u}_b$  is related to  $w$  by (2.26). By the same argument as above, for every small  $b \in \mathbb{R}$  and  $\delta \in \mathbb{R}$ , there exists a unique solution  $w \in H^2_{\text{even}}$  of equation (2.30) satisfying the bound

$$(2.31) \quad \|w\|_{H^2} \leq A(\delta^2 + |\delta||b| + |b|^3),$$

where the positive constant  $A$  is independent of  $\delta$  and  $b$ .

Substituting the near-identity transformation (2.26) into the bifurcation equation (2.24) and using the bound (2.31) for the component  $w \in H^2_{\text{even}}$ , we rewrite (2.24) in the equivalent form

$$(2.32) \quad \alpha \delta b + \beta b^3 + \mathcal{O}(\delta b^2, b^4) = 0,$$

where we have introduced numerical coefficients  $\alpha$  and  $\beta$  as follows:

$$(2.33) \quad \alpha := -4\|\psi_*\|_{L^2}^2 + 12\langle \psi_*^2, \partial_c u_{c_*} \rangle_{L^2}$$

and

$$(2.34) \quad \beta := 12\langle \psi_*^2, w_0 + w_2 \rangle_{L^2}.$$



We have also removed  $\mathcal{O}(\delta^2)$  from the remainder term in (2.32), because the bifurcation equation (2.24) is identically satisfied in the case of the line soliton with  $b = 0$  for every small  $\delta \in \mathbb{R}$ .

Comparison of (2.27) and (2.33) with (2.15) shows that

$$\alpha \equiv -\langle \psi_*, L'_{c_*} \psi_* \rangle_{L^2} = \frac{16}{3\sqrt{c_*}}.$$

On the other hand, the coefficient  $\beta$  in (2.34) is less explicit. In order to show that  $\beta < 0$ , we obtain the unique even solution of the linear inhomogeneous equation (2.28) in the explicit form

$$(2.35) \quad w_0(\xi) = -15\operatorname{sech}^2(\sqrt{c_*}\xi) + \frac{15}{2}\operatorname{sech}^4(\sqrt{c_*}\xi),$$

where we have used  $c_* = \frac{1}{5}$ . Because  $L_{c_*} + 4 : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is strictly positive, there exists a unique solution of the linear inhomogeneous equation (2.29). Unfortunately, it is not available in the explicit form. Nevertheless, we can represent this unique solution of equation (2.29) in the form

$$(2.36) \quad w_2(\xi) = 5\operatorname{sech}^2(\sqrt{c_*}\xi) + \frac{15}{4}\operatorname{sech}^4(\sqrt{c_*}\xi) - \tilde{w}_2(\xi),$$

where  $\tilde{w}_2$  is found from the inhomogeneous equation

$$(2.37) \quad (L_{c_*} + 4)\tilde{w}_2(\xi) = 20\operatorname{sech}^2(\sqrt{c_*}\xi).$$

By the maximum principle for the elliptic operator  $(L_{c_*} + 4)$ , the component  $\tilde{w}_2$  satisfies  $\tilde{w}_2(\xi) \geq 0$  for all  $\xi \in \mathbb{R}$ . After computing the integrals in (2.34), we obtain

$$(2.38) \quad \beta = 12\langle \psi_*^2, w_0 + w_2 \rangle_{L^2} = -12\langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} < 0.$$

The cubic equation (2.22) follows from the truncation of the bifurcation equation (2.32) at the first two terms. This is justified because if  $b \neq 0$ , then it follows from (2.32) that

$$\delta = -\frac{\beta}{\alpha}b^2 + \mathcal{O}(b^3)$$

and if  $b_0$  is a root of the cubic equation (2.22) for a given small  $\delta = c - c_*$ , then

$$|b - b_0| \leq Ab_0^2,$$

where the positive constant  $A$  is independent of  $\delta$ . Then, it follows from the bound (2.25) that  $\|\tilde{u}_b\|_{H^2} \leq Ab_0^2$ , which justifies the decomposition (2.21) after the change of notation  $b_0 \mapsto b$ .  $\blacksquare$

**Remark 2.5.** *Since  $\alpha > 0$  and  $\beta < 0$ , the nonzero solutions for  $b$  exists in the cubic equation (2.22) if and only if  $c > c_*$ . By Lemma 2.1, the line soliton with  $c > c_*$  is spectrally unstable with respect to the transverse  $2\pi$ -periodic perturbations.*

**2.4. Statement of the main theorem.** The ZK equation (1.1) was shown in [12] to be locally well-posed in  $H^s(\mathbb{R} \times \mathbb{T})$  for  $s > \frac{3}{2}$  and globally well-posed for perturbations of the line solitons (2.1) in  $H^1(\mathbb{R}^2)$ . More recently, the ZK equation (1.1) was shown in [14] to be globally well-posed in  $H^1(\mathbb{R} \times \mathbb{T})$ . The latter well-posedness result allows us to employ the energy method in the justification of the normal form for transverse instability of the line soliton with a nearly critical speed of propagation.

Let us denote by  $H_\mu^s(\mathbb{R} \times \mathbb{T})$  the exponentially weighted version of the space  $H^s(\mathbb{R} \times \mathbb{T})$ ,  $s \geq 0$  with the weight  $\mu > 0$  applied in the  $\xi$  axis only.

In order to characterize the dynamics of solutions of the ZK equation (1.1) in time  $t$  near the line solitons (2.1), we introduce varying parameters  $a(t)$  and  $c(t)$  of the line solitons as well as its perturbation  $\tilde{u}(t)$  defined in  $H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T})$  for every  $t \in \mathbb{R}_+$ . Hence, we introduce the travelling coordinate  $\xi = x - 4a(t)$  and use the decomposition

$$(2.39) \quad u(x, y, t) = u_{c(t)}(\xi) + \tilde{u}(\xi, y, t), \quad \xi = x - 4a(t).$$

The time evolution of the varying parameters  $a(t)$  and  $c(t)$  and the perturbation term  $\tilde{u}(t)$  are to be found from the evolution problem

$$(2.40) \quad \tilde{u}_t = \partial_\xi(L_c - \partial_y^2 + 4(\dot{a} - c))\tilde{u} + 4(\dot{a} - c)\partial_\xi u_c - \dot{c}\partial_c u_c - 6\partial_\xi \tilde{u}^2,$$

where the differential expression for  $L_c$  is given by (2.3).

For  $c = c_*$ , the operators  $L_{c_*} + k^2 : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  are coercive for any  $k \in \mathbb{Z} \setminus \{0, \pm 1\}$ . This property will be used to control perturbations with the corresponding Fourier wave numbers. On the other hand, the operator  $\partial_\xi L_{c_*} : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  has a double zero eigenvalue associated with the Jordan block (2.8), whereas the operator  $\partial_\xi(L_{c_*} + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  has a simple zero eigenvalue by Lemma 2.2. The double zero eigenvalue of  $\partial_\xi L_{c_*} : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  is already incorporated in the decomposition (2.39), whereas the simple zero eigenvalue of  $\partial_\xi(L_{c_*} + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  will be used in the secondary decomposition of the perturbation term  $\tilde{u}$ .

The following theorem represents the normal form for transverse instability of the line soliton with a nearly critical speed of propagation.

**Theorem 1.** *Consider the Cauchy problem for the evolution equation (2.40) with*

$$\tilde{u}(0) \in H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}),$$

where  $\mu > 0$  is sufficiently small. There exist  $\varepsilon_0 > 0$  and  $C_0 > 0$  such that if the initial data satisfy the bound

$$(2.41) \quad \|\tilde{u}(0) - 2\varepsilon \cos(y)\psi_*\|_{H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T})} + |c(0) - c_*| \leq \varepsilon^2,$$

for every  $\varepsilon \in (0, \varepsilon_0)$ , then there exist unique functions  $a, b, c \in C^1(\mathbb{R}_+)$  and the unique solution

$$\tilde{u}(t) \in C(\mathbb{R}_+; H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}))$$

of the evolution equation (2.40) satisfying the bound

$$(2.42) \quad \|\tilde{u}(t) - (b(t)e^{iy} + \bar{b}(t)e^{-iy})\psi_*\|_{H^1(\mathbb{R} \times \mathbb{T})} + |c(t) - c_*| \leq C_0\varepsilon^2, \quad t \in \mathbb{R}_+,$$

Furthermore, there exists  $c_+ \in \mathbb{R}$  such that the function  $b(t)$  satisfies the normal form equation

$$(2.43) \quad \dot{b} = \lambda'(c_*)(c_+ - c_*)b + \gamma|b|^2b, \quad t \in \mathbb{R}_+,$$

with  $b(0) = \varepsilon$  and  $|c_+ - c_*| \leq C_0 \varepsilon^2$ , where  $\lambda'(c_*) > 0$  is given by (2.13) and  $\gamma < 0$  is a specific numerical coefficient given by (3.31) below. Consequently,  $|b(t)| \leq C_0 \varepsilon$  for every  $t \in \mathbb{R}_+$ .

**Remark 2.6.** *The linear part of the normal form (2.43) reproduces the spectral stability result of Lemma 2.2.*

**Remark 2.7.** *The stationary part of the normal form (2.43) represents the bifurcation result (2.22) in Lemma 2.3, except for the quantitative discrepancy between the numerical coefficients  $(\alpha, \beta)$  and numerical coefficients  $(\lambda'(c_*), \gamma)$ , see Remark 3.1 below.*

**Remark 2.8.** *Similarly to the asymptotic stability result in [17], we anticipate that for  $\mu > 0$  sufficiently small, there exists  $b_\infty$  and  $c_\infty$  such that the solution in Theorem 1 satisfies the following limits:*

$$(2.44) \quad c_+ \leq c_* : \quad \lim_{t \rightarrow \infty} \|u(t) - u_{c_\infty}\|_{H_\mu^1(\mathbb{R} \times \mathbb{T})} = 0,$$

and

$$(2.45) \quad c_+ > c_* : \quad \lim_{t \rightarrow \infty} \|u(t) - u_{b_\infty}\|_{H_\mu^1(\mathbb{R} \times \mathbb{T})} = 0,$$

where  $u_{c_\infty}$  is the line soliton (2.1) with  $c = c_\infty$  and  $u_{b_\infty}$  is the transversely modulated solitary wave defined by Lemma 2.2 with  $b = b_\infty$ . The limits (2.44) and (2.45) are in agreement with the asymptotic stability results obtained in [27]. However, the proof of these limits requires more control of the modulation equations for perturbations in  $H_\mu^1(\mathbb{R} \times \mathbb{T})$ . The relevant tools are not available from the previous work [17], where perturbations were considered in  $H_\mu^1(\mathbb{R})$ .

### 3. DERIVATION AND JUSTIFICATION OF THE NORMAL FORM (2.43)

We first derive the general modulation equations for the varying parameters  $a$  and  $c$  in the decomposition (2.39) near the line solitons (2.1). Next, we introduce the varying parameter  $b$  in the secondary decomposition along the neutral eigenmode  $\psi_*$  in Lemma 2.2 and derive the corresponding modulation equation for  $b$ . Further, we simplify the modulation equations and derive the normal form (2.43) by means of nearly identity transformations. Finally, we justify the bound (2.42) with the energy method.

**3.1. Modulation equations for the varying parameters  $a$  and  $c$ .** Let  $X_c = \text{span}\{\partial_\xi u_c, \partial_c u_c\}$  be an invariant subspace of  $L^2(\mathbb{R})$  for the double zero eigenvalue of the linearized operator  $\partial_\xi L_c : H^3(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ , according to the Jordan block (2.8). Thanks to the exponential decay of  $u_c(\xi)$  as  $|\xi| \rightarrow \infty$ , there is  $\mu_0 > 0$  such that  $X_c$  is also an invariant subspace of  $L_\mu^2(\mathbb{R})$  for  $\partial_\xi L_c : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$  for  $\mu \in (0, \mu_0)$ . Similarly,  $X_c^* = \text{span}\{u_c, \partial_\xi^{-1} \partial_c u_c\}$  is an invariant subspace of  $L_{-\mu}^2(\mathbb{R})$  for the double zero eigenvalue of the adjoint operator  $-L_c \partial_\xi : H_{-\mu}^3(\mathbb{R}) \rightarrow L_{-\mu}^2(\mathbb{R})$ . Recall that the exponentially weighted space is defined in (2.9) and

$$\partial_\xi^{-1} u(\xi) := \int_{-\infty}^{\xi} u(\xi') d\xi'.$$

In order to avoid confusion between spaces  $L_\mu^2(\mathbb{R})$  and  $L_\mu^2(\mathbb{R} \times \mathbb{T})$ , we specify whether the spatial domain is  $\mathbb{R}$  or  $\mathbb{R} \times \mathbb{T}$  in the  $L^2$  inner products and their induced norms.

The following lemma states the validity of the decomposition (2.39).

**Lemma 3.1.** *There exists  $\varepsilon_0 > 0$ ,  $\mu_0 > 0$ , and  $C_0 > 0$  such that if  $u \in C(\mathbb{R}_+, H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}))$  with  $\mu \in (0, \mu_0)$  is a global solution to the ZK equation (1.1) satisfying*

$$(3.1) \quad \varepsilon := \inf_{a \in \mathbb{R}} \|u(x + 4a, y, t) - u_{c_*}(x)\|_{H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T})} \leq \varepsilon_0, \quad t \in \mathbb{R}_+,$$

*then there exist  $a, c \in C(\mathbb{R}_+)$  and  $\tilde{u} \in C(\mathbb{R}_+, H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}))$  such that the decomposition*

$$(3.2) \quad u(x, y, t) = u_{c(t)}(\xi) + \tilde{u}(\xi, y, t), \quad \xi = x - 4a(t)$$

*holds with  $\tilde{u}(t) \in [X_{c(t)}^*]^\perp$  for every  $t \in \mathbb{R}_+$ , where*

$$(3.3) \quad [X_{c(t)}^*]^\perp = \{\tilde{u} \in L_\mu^2(\mathbb{R} \times \mathbb{T}) : \langle u_{c(t)}, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = \langle \partial_\xi^{-1} \partial_c u_{c(t)}, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = 0\}.$$

*Moreover,  $c(t)$  and  $\tilde{u}(t)$  satisfies*

$$(3.4) \quad |c(t) - c_*| + \|\tilde{u}(t)\|_{H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T})} \leq C\varepsilon, \quad t \in \mathbb{R}_+.$$

**Proof.** The proof is relatively well-known, see Proposition 5.1 in [17]. It is based on the implicit function theorem applied to the two constraints in the definition of  $[X_{c(t)}^*]^\perp$  in (3.3).  $\blacksquare$

By the global well-posedness theory for the ZK equation (1.1) [14], there exists a unique global solution in class  $u \in C(\mathbb{R}_+, H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}))$  for  $\mu > 0$  sufficiently small. By the initial bound (2.41), the initial data satisfy (3.1) for some  $\varepsilon > 0$  sufficiently small. By the elementary continuation arguments, the decomposition (3.2) can be used as long as the solution  $u$  satisfies (3.1). The component  $\tilde{u}$  in the decomposition (3.2) satisfies the evolution equation (2.40) rewritten again as

$$(3.5) \quad \tilde{u}_t = \partial_\xi(L_c - \partial_y^2 + 4(\dot{a} - c))\tilde{u} + 4(\dot{a} - c)\partial_\xi u_c - \dot{c}\partial_c u_c - 6\partial_\xi \tilde{u}^2,$$

where the differential expression for  $L_c$  is given by (2.3).

Both parameters  $a$  and  $c$  depend on the time variable  $t$ . Modulation equations for  $a$  and  $c$  are derived from the well-known projection algorithm, which has been applied to similar problems in [2, 13, 17]. The two constraints on  $\tilde{u}$  in (3.3) represent the symplectic orthogonality conditions, which specify uniquely  $a$  and  $c$  in the decomposition (3.2) as well as the time evolution of  $a$  and  $c$ . Moreover, one can show that  $a, c \in C^1(\mathbb{R}_+)$ .

If  $\tilde{u}(0) \in [X_{c(0)}^*]^\perp$  initially, then  $\tilde{u}(t)$  remains in  $[X_{c(t)}^*]^\perp$  for every  $t \in \mathbb{R}_+$ , provided that the varying parameters  $a, c \in C^1(\mathbb{R}_+)$  satisfy the system of two modulation equations

$$(3.6) \quad \begin{bmatrix} \frac{1}{2}(M'(c))^2 - \frac{1}{2\pi} \langle \partial_\xi^{-1} \partial_c^2 u_c, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} & P'(c) + \frac{1}{2\pi} \langle \partial_c u_c, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \\ P'(c) - \frac{1}{2\pi} \langle \partial_c u_c, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} & \frac{1}{2\pi} \langle \partial_\xi u_c, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \end{bmatrix} \begin{bmatrix} \dot{c} \\ 4(\dot{a} - c) \end{bmatrix} = \frac{3}{\pi} \begin{bmatrix} \langle \partial_c u_c, \tilde{u}^2 \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \\ \langle \partial_\xi u_c, \tilde{u}^2 \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \end{bmatrix},$$

where  $M(c) = \int_{\mathbb{R}} u_c(\xi) d\xi$  and  $P(c) = \frac{1}{2} \int_{\mathbb{R}} u_c^2(\xi) d\xi$ . From the expression (2.1), we obtain  $M'(c) = 1/\sqrt{c}$  and  $P'(c) = \sqrt{c}$ .

**3.2. A secondary decomposition for  $c = c_*$ .** By Lemma 2.2, if  $c = c_* = \frac{1}{5}$  and  $\mu > 0$  is sufficiently small, then  $Y_{c_*} = \text{span}\{\psi_*\}$  is an invariant subspace of  $L_\mu^2(\mathbb{R})$  for the simple zero eigenvalue of the linearized operator  $\partial_\xi(L_{c_*} + 1) : H_\mu^3(\mathbb{R}) \rightarrow L_\mu^2(\mathbb{R})$ . Similarly,  $Y_{c_*}^* = \text{span}\{\eta_*\}$  is an invariant subspace of  $L_{-\mu}^2(\mathbb{R})$  for the simple zero eigenvalue of the adjoint operator  $-(L_{c_*} + 1)\partial_\xi : H_{-\mu}^3(\mathbb{R}) \rightarrow L_{-\mu}^2(\mathbb{R})$ . We note the double degeneracy of the Fourier harmonics  $e^{iy}$  and  $e^{-iy}$ , when general transverse perturbations are considered.

The following lemma states the secondary decomposition of the solution  $\tilde{u}$  defined in the primary decomposition (3.2).

**Lemma 3.2.** *Under assumptions of Lemma 3.1, let  $\tilde{u} \in C(\mathbb{R}_+, H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}))$  be given by the decomposition (3.2) and (3.3). There exist  $b \in C(\mathbb{R}_+)$  and  $v \in C(\mathbb{R}_+, H^1(\mathbb{R} \times \mathbb{T}) \cap H_\mu^1(\mathbb{R} \times \mathbb{T}))$  such that the decomposition*

$$(3.7) \quad \tilde{u}(\xi, y, t) = (b(t)e^{iy} + \bar{b}(t)e^{-iy})\psi_*(\xi) + v(\xi, y, t),$$

holds with  $v(t) \in [Y_{c(t)}^*]^\perp$  for every  $t \in \mathbb{R}_+$ , where

$$(3.8) \quad [Y_{c(t)}^*]^\perp = \{v \in [X_{c(t)}^*]^\perp : \langle \eta_* e^{iy}, v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = \langle \eta_* e^{-iy}, v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = 0\}.$$

**Proof.** The proof is straightforward thanks to the fact  $\langle \eta_*, \psi_* \rangle_{L^2(\mathbb{R})} \neq 0$  by (2.11).  $\blacksquare$

We introduce the decomposition

$$(3.9) \quad a(t) = \int_0^t c(t') dt' + h(t), \quad c(t) = c_* + \delta(t),$$

in addition to the decomposition (3.7). To simplify notations, we also write  $L_c = L_{c_*} + \Delta L_c$ , where  $\Delta L_c \in L^\infty(\mathbb{R})$  satisfies the bound  $\|\Delta L_c\|_{L^\infty} \leq A|c - c_*|$  for  $|c - c_*|$  sufficiently small with a  $c$ -independent positive constant  $A$ .

The correction term  $v$  in the decomposition (3.7) satisfies the time evolution equation

$$(3.10) \quad \begin{aligned} v_t &= \partial_\xi(L_{c_*} - \partial_y^2 + 4\dot{h} + \Delta L_c)v + 4\dot{h}\partial_\xi u_{c_*+\delta} - \dot{\delta}\partial_c u_{c_*+\delta} - (\dot{b}e^{iy} + \dot{\bar{b}}e^{-iy})\psi_* \\ &+ \partial_\xi(4\dot{h} + \Delta L_c)(be^{iy} + \bar{b}e^{-iy})\psi_* - 6\partial_\xi((be^{iy} + \bar{b}e^{-iy})\psi_* + v)^2. \end{aligned}$$

The two constraints in (3.8) represent the symplectic orthogonality conditions, which specify uniquely the complex parameter  $b$  in the secondary decomposition (3.7). Again, one can show that  $b \in C^1(\mathbb{R}_+)$

If  $v(0) \in [Y_{c(0)}^*]^\perp$  initially, then  $v(t)$  remains in  $[Y_{c(t)}^*]^\perp$  for every  $t \in \mathbb{R}_+$ , provided that the varying parameter  $b \in C^1(\mathbb{R}_+)$  satisfy the following modulation equation:

$$(3.11) \quad \begin{aligned} &\dot{b}\langle \eta_*, \psi_* \rangle_{L^2(\mathbb{R})} + b\langle \psi_*, (4\dot{h} + \Delta L_c)\psi_* \rangle_{L^2(\mathbb{R})} + \frac{1}{2\pi}\langle \psi_* e^{iy}, \Delta L_c v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \\ &= \frac{3}{\pi}\langle \psi_* e^{iy}, [(be^{iy} + \bar{b}e^{-iy})\psi_* + v]^2 \rangle_{L^2(\mathbb{R} \times \mathbb{T})}. \end{aligned}$$

Substituting (3.7) and (3.9) to the system (3.6) yields the equivalent form of the modulation equations:

$$(3.12) \quad \begin{aligned} & \left[ \begin{array}{cc} \frac{1}{2}(M'(c))^2 - \frac{1}{2\pi} \langle \partial_\xi^{-1} \partial_c^2 u_c, v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} & P'(c) + \frac{1}{2\pi} \langle \partial_c u_c, v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \\ P'(c) - \frac{1}{2\pi} \langle \partial_c u_c, v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} & \frac{1}{2\pi} \langle \partial_\xi u_c, v \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \end{array} \right] \begin{bmatrix} \dot{\delta} \\ 4\dot{h} \end{bmatrix} \\ &= \frac{3}{\pi} \begin{bmatrix} \langle \partial_c u_c, [(be^{iy} + \bar{b}e^{-iy})\psi_* + v]^2 \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \\ \langle \partial_\xi u_c, [(be^{iy} + \bar{b}e^{-iy})\psi_* + v]^2 \rangle_{L^2(\mathbb{R} \times \mathbb{T})} \end{bmatrix}, \end{aligned}$$

where  $c(t) = c_* + \delta(t)$ . The system (3.11) and (3.12) determine the time evolution of the varying parameters  $b$ ,  $h$ , and  $\delta$ , whereas the evolution problem (3.10) determines the correction term  $v(t) \in [Y_{c(t)}^*]^\perp$ .

**3.3. Near-identity transformations.** Because  $u_{c_*}$  and  $\psi_*^2$  are even functions of  $\xi$ , whereas  $P'(c_*) \neq 0$ , the modulation equations (3.11) and (3.12) yields the following balance at the leading order:

$$\dot{b} = \mathcal{O}((|\delta| + |b|^2)|b|), \quad \dot{h} = \mathcal{O}(|b|^2),$$

whereas the source terms in the evolution problem (3.10) are of the order of  $\mathcal{O}(|b|^2)$ . This suggests the use of near-identity transformation, which mimics the ones used in the proof of Lemma 2.3. In particular, we will remove the  $\mathcal{O}(|b|^2)$  terms in the equation for  $\dot{h}$  and  $v_t$ . Hence, we represent the correction term  $v$  in the decomposition (3.7) as follows:

$$(3.13) \quad v(\xi, y, t) = (b(t)^2 e^{2iy} + \bar{b}(t)^2 e^{-2iy}) w_2(\xi) + |b(t)|^2 w_0(\xi) + w(\xi, y, t),$$

where  $w_0$  and  $w_2$  are the same solutions of the linear inhomogeneous equations (2.28) and (2.29), whereas  $w$  satisfies the transformed evolution equation

$$(3.14) \quad \begin{aligned} w_t &= \partial_\xi(L_{c_*} - \partial_y^2 + 4\dot{h} + \Delta L_c)w + 4\dot{h}\partial_\xi u_{c_*+\delta} - \dot{\delta}\partial_c u_{c_*+\delta} \\ &\quad - (\dot{b}e^{iy} + \dot{\bar{b}}e^{-iy})\psi_* - (2\dot{b}\dot{b}e^{2iy} + 2\dot{\bar{b}}\dot{\bar{b}}e^{-2iy})w_2 - (\dot{\bar{b}}\dot{b} + \dot{b}\dot{\bar{b}})w_0 \\ &\quad + \partial_\xi \left( 4\dot{h} + \Delta L_c \right) [(be^{iy} + \bar{b}e^{-iy})\psi_* + (b^2 e^{2iy} + \bar{b}^2 e^{-2iy})w_2 + |b|^2 w_0] \\ &\quad - 12\partial_\xi (be^{iy} + \bar{b}e^{-iy})\psi_* [(b^2 e^{2iy} + \bar{b}^2 e^{-2iy})w_2 + |b|^2 w_0] \\ &\quad - 6\partial_\xi [(b^2 e^{2iy} + \bar{b}^2 e^{-2iy})w_2 + |b|^2 w_0]^2. \end{aligned}$$

We rewrite the first equation in the system (3.12) as follows:

$$(3.15) \quad 4P'(c_*)\dot{h} = 12|b|^2 \langle \partial_c u_{c_*}, \psi_*^2 \rangle_{L^2} + \mathcal{O}(|b|^4) = \frac{48}{5\sqrt{c_*}}|b|^2 + \mathcal{O}(|b|^4),$$

where the explicit expression (2.27) has been used. Since  $P'(c_*) = \sqrt{c_*}$  and  $c_* = \frac{1}{5}$ , we obtain

$$(3.16) \quad \dot{h} = 12|b|^2 + \mathcal{O}(|b|^4)$$

and

$$(3.17) \quad w(\xi, y, t) = 12|b|^2 \partial_c u_{c_*} + \tilde{w}(\xi, y, t),$$

where  $\tilde{w}$  satisfied a transformed evolution equation without the  $\mathcal{O}(|b|^2)$  terms in the right-hand side of (3.14). Substituting (3.13), (3.16), and (3.17) into the modulation equation

(3.11) yields

$$(3.18) \quad \begin{aligned} \dot{b}\langle \eta_*, \psi_* \rangle_{L^2} &= 12|b|^2 b \langle \psi_*^2, w_0 + w_2 \rangle_{L^2} + 144|b|^2 b \langle \psi_*^2, \partial_c u_{c_*} \rangle_{L^2} \\ &\quad - b \langle \psi_*, (L'_{c_*} \delta + 48|b|^2) \psi_* \rangle_{L^2} + \mathcal{O}(\delta^2 |b| + |b|^5), \end{aligned}$$

where  $L'_{c_*}$  is given by (2.15) and we have used  $\Delta L_c = L'_{c_*} \delta + \mathcal{O}(\delta^2)$  for  $\delta = c - c_*$ . After straightforward computations, equation (3.18) takes the form

$$(3.19) \quad \dot{b}\langle \eta_*, \psi_* \rangle_{L^2} = -12|b|^2 b \langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} + \frac{64}{\sqrt{c_*}} |b|^2 b + \frac{16}{3\sqrt{c_*}} b \delta + \mathcal{O}(\delta^2 |b| + |b|^5),$$

where  $\tilde{w}_2$  is found from the solution of the linear inhomogeneous equation (2.37).

The normal form (3.19) is not closed because  $\delta$  is related to  $|b|^2$  by the second equation of the system (3.12). In fact, this equation relates  $\dot{\delta}$  to  $\bar{b}\dot{b} + \dot{b}b$ , however, it yields  $\delta = \mathcal{O}(|b|^2)$  after integration. We write the second equation in the system (3.12) as follows:

$$(3.20) \quad P'(c_*) \dot{\delta} = \frac{6}{\pi} \langle \psi_* \partial_\xi u_{c_*}, (be^{iy} + \bar{b}e^{-iy}) \tilde{w} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} + \mathcal{O}(|b|^3 |\dot{b}|).$$

Note that many  $\mathcal{O}(|b|^4)$  terms in the second equation in the system (3.12) are identically zero because the  $L^2$  scalar products are computed between odd and even functions. We are now looking at the  $\mathcal{O}(|\dot{b}|)$  terms in  $\tilde{w}$ , which contributes to the leading order of equation (3.20). Substituting (3.17) into (3.14) yields the following evolution equation for  $\tilde{w}$ :

$$(3.21) \quad \begin{aligned} \tilde{w}_t &= \partial_\xi (L_{c_*} - \partial_y^2 + 4\dot{h} + \Delta L_c) \tilde{w} - (\dot{b}e^{iy} + \dot{\bar{b}}e^{-iy}) \psi_* + \partial_\xi \left( \frac{48}{5c_*} |b|^2 + \delta L'_{c_*} \right) (be^{iy} + \bar{b}e^{-iy}) \psi_* \\ &\quad - 12\partial_\xi (be^{iy} + \bar{b}e^{-iy}) \psi_* [(b^2 e^{2iy} + \bar{b}^2 e^{-2iy}) w_2 + |b|^2 w_0] + \mathcal{O}(|b|^4). \end{aligned}$$

The terms proportional to  $\dot{b}$  have different parity in spatial coordinate  $\xi$  compared to the terms proportional to  $|b|^2 b$  and  $\delta b$ . Only odd terms in  $\xi$  give a nonzero contribution to equation (3.20). Therefore, we single out the odd terms in  $\xi$  from the other terms in the representation:

$$(3.22) \quad \tilde{w}(\xi, y, t) = \left( \dot{b}(t) e^{iy} + \dot{\bar{b}}(t) e^{-iy} \right) w_3(\xi) + \tilde{\tilde{w}}(\xi, y, t),$$

where  $w_3 \in L^2_\mu(\mathbb{R})$  is a solution of the linear inhomogeneous equation

$$(3.23) \quad \partial_\xi (L_{c_*} + 1) w_3 = \psi_* + \frac{\langle \eta_*, \psi_* \rangle_{L^2}}{\|\psi_*\|_{L^2}^2} \partial_\xi \psi_*,$$

whereas  $\tilde{\tilde{w}}$  satisfies a transformed equation without the  $\dot{b}$  terms in the right-hand side of (3.21). The additional projection term is added to the right-hand-side of equation (3.23) to ensure that the Fredholm's solvability condition is satisfied for this inhomogeneous equation in  $L^2_\mu(\mathbb{R})$ . Integrating equation (3.23) yields the inhomogeneous equation for  $w_3 \in L^2_\mu(\mathbb{R})$ :

$$(3.24) \quad (L_{c_*} + 1) w_3 = \int_{+\infty}^\xi \psi_*(\xi') d\xi' + \frac{\langle \eta_*, \psi_* \rangle_{L^2}}{\|\psi_*\|_{L^2}^2} \psi_*,$$

where the limits of integrations ensure the correct decay as  $\xi \rightarrow +\infty$  in (2.9).

Let  $w_4 \in H^2(\mathbb{R})$  be a unique odd solution of the linear inhomogeneous equation

$$(3.25) \quad (L_{c_*} + 1) w_4 = \psi_* \partial_\xi u_{c_*}.$$

The unique solution of equation (3.25) exists in the explicit analytical form

$$(3.26) \quad w_4(\xi) = -\frac{\sqrt{c_*}}{4} \tanh(\sqrt{c_*}\xi) \operatorname{sech}^3(\sqrt{c_*}\xi) = \frac{1}{12} \partial_\xi \psi_*(\xi).$$

By using (3.24) and (3.26), we compute

$$(3.27) \quad \langle \psi_* \partial_\xi u_{c_*}, w_3 \rangle_{L^2} = \langle (L_{c_*} + 1)w_4, w_3 \rangle_{L^2} = \langle w_4, (L_{c_*} + 1)w_3 \rangle_{L^2} = -\frac{1}{12} \|\psi_*\|_{L^2}^2 = -\frac{4}{45\sqrt{c_*}}.$$

Equation (3.20) with (3.27) and  $P'(c_*) = \sqrt{c_*}$  yields

$$(3.28) \quad \frac{d\delta}{dt} = \frac{12}{\sqrt{c_*}} \frac{d|b|^2}{dt} \langle \psi_* \partial_\xi u_{c_*}, w_3 \rangle_{L^2} + \mathcal{O}(|b|^3|\dot{b}|) = -\frac{16}{3} \frac{d|b|^2}{dt} + \mathcal{O}(|b|^3|\dot{b}|).$$

Integrating this equation in time, we finally obtain

$$(3.29) \quad \delta(t) = \delta_0 - \frac{16}{3} |b(t)|^2 + \mathcal{O}(|b(t)|^4),$$

where  $\delta_0$  is an arbitrary integration constant. Substituting equation (3.29) into the normal form (3.19) yield

$$(3.30) \quad \dot{b} \langle \eta_*, \psi_* \rangle_{L^2} = -12|b|^2 b \langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} + \frac{16}{3\sqrt{c_*}} \left( \delta_0 + \frac{20}{3} |b|^2 \right) b + \mathcal{O}(\delta_0^2 |b| + \delta_0 |b|^3 + |b|^5).$$

Defining  $\delta_0 := c_+ - c_*$ , using the explicit expression (2.11) and (2.13), and truncating (3.30) yield the normal form (2.43) with

$$(3.31) \quad \gamma := \frac{96}{\pi^2 \sqrt{c_*}} \left( -\frac{1}{5} \sqrt{c_*} \langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} + \frac{16}{27} \right).$$

Under the assumption that  $|\delta_0| \leq C_0 \varepsilon^2$  for some  $\varepsilon$ -independent positive constant  $C_0$ , the remainder terms in (3.30) is of the order  $\mathcal{O}(\varepsilon^5)$ , which does not contribute to the leading order of the bound (2.42). Therefore, the truncation of (3.30) into (2.43) is justified within the approximation error in (2.42).

**Remark 3.1.** *The numerical coefficients  $(\lambda(c_*), \gamma)$  are different from the numerical coefficients  $(\alpha, \beta)$ . This difference is explained as follows. Expansions (3.9), (3.16), and (3.29) yield*

$$c = c_* + \delta_0 - \frac{16}{3} |b|^2 + \mathcal{O}(|b|^4), \quad \dot{a} = c_* + \delta_0 + \frac{20}{3} |b|^2 + \mathcal{O}(|b|^4).$$

*The normal form (3.30) in the stationary case  $\dot{b} = 0$  corresponds to the effective speed correction given by*

$$\dot{a} - c_* = \delta_0 + \frac{20}{3} |b|^2 + \mathcal{O}(|b|^4) = \frac{9}{4} \sqrt{c_*} |b|^2 \langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} + \mathcal{O}(|b|^4) = -\frac{\beta}{\alpha} |b|^2 + \mathcal{O}(|b|^4),$$

*in agreement with the cubic algebraic equation (2.22). In the time-dependent case, the roles of  $\delta_0$  and  $\frac{20}{3} |b|^2$  are different because the former is constant in  $t$  but the latter changes in  $t$ .*

Bifurcation analysis of Lemma 2.3 relies on the fact that the coefficient  $\beta$  of the cubic term in the normal form (2.22) is negative. It is equally important for the stability analysis near the line soliton that the coefficient  $\gamma$  of the cubic term in the normal form (2.43) is negative. Since  $\tilde{w}_2(\xi) \geq 0$  for all  $\xi \in \mathbb{R}$ , the first term in  $\gamma$  is negative. On the other hand,



the second term is positive, so that  $\gamma < 0$  if the negative term prevails. Let us show it numerically.

We approximate the function  $\tilde{w}_2$  numerically by using the central-difference method for the linear inhomogeneous equation (2.37) and then approximate the integral  $\langle \psi_*^2, \tilde{w}_2 \rangle_{L^2}$  by using the composite trapezoidal method. Testing the codes on the function  $w_0$  which satisfies the linear inhomogeneous equation (2.28) with the explicit solution (2.35), we obtain numerically that

$$\frac{1}{5} \sqrt{c_*} \langle \psi_*^2, w_0 \rangle_{L^2} \approx -1.5238$$

which corresponds to the exact value  $-\frac{32}{21}$  within the computational error of  $\mathcal{O}(10^{-5})$ . Performing the same task for  $\tilde{w}_2$  which satisfies the linear inhomogeneous equation (2.37), we obtain

$$\frac{1}{5} \sqrt{c_*} \langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} \approx 1.2359,$$

which is essentially bigger than  $\frac{16}{27} \approx 0.5926$ . Therefore,

$$\frac{1}{5} \sqrt{c_*} \langle \psi_*^2, \tilde{w}_2 \rangle_{L^2} > \frac{16}{27},$$

which implies that  $\gamma < 0$  in (3.31).

**3.4. Justification of the approximation error.** It remains to justify the error bound (2.42). We achieve it with the energy method pioneered in [17]. First, we recall that the energy

$$(3.32) \quad E(u) = \frac{1}{2} \int_{\mathbb{R} \times \mathbb{T}} [u_x^2 + u_y^2 - 4u^3] dx dy$$

and the momentum

$$(3.33) \quad P(u) = \frac{1}{2} \int_{\mathbb{R} \times \mathbb{T}} u^2 dx dy$$

are conserved in time  $t$  for a global solution  $u \in C(\mathbb{R}, H^1(\mathbb{R} \times \mathbb{T}))$  to the ZK equation (1.1). The line soliton (2.1) is a critical point of the action functional  $\Lambda_c(u) := E(u) + 4cP(u)$ , see (2.2). Thanks to the translational invariance of the ZK equation (1.1), the decomposition (2.39) yields

$$(3.34) \quad \Lambda_c(u_c + \tilde{u}) - \Lambda_c(u_c) = \frac{1}{2} \langle (L_c - \partial_y^2) \tilde{u}, \tilde{u} \rangle_{L^2} + N_c(\tilde{u}),$$

where the differential expression for  $L_c$  is given by (2.3) and  $N_c$  is a nonlinear term satisfying

$$(3.35) \quad |N_c(\tilde{u})| \leq A \|\tilde{u}\|_{H^1}^3,$$

for some positive constant  $A$  as long as  $\|\tilde{u}\|_{H^1}$  is small. By using the Fourier series

$$\tilde{u}(\xi, y, t) = \frac{1}{\sqrt{2\pi}} \sum_{k \in \mathbb{Z}} \hat{u}_k(\xi, t) e^{iky}$$

and Parseval's equality, we can represent the second variation of  $\Lambda_c$  at  $u_c$  in the form

$$(3.36) \quad \langle (L_c - \partial_y^2) \tilde{u}, \tilde{u} \rangle_{L^2(\mathbb{R} \times \mathbb{T})} = \sum_{k \in \mathbb{Z}} \langle (L_c + k^2) \hat{u}_k, \hat{u}_k \rangle_{L^2(\mathbb{R})}.$$

The following lemma summarizes the coercivity results for the second variation of  $\Lambda_c$  at  $u_c$ .

**Lemma 3.3.** *For  $\mu > 0$  sufficiently small, there exists a constant  $A > 0$  such that for every  $\hat{u}_k \in H^1(\mathbb{R})$  and for every  $k \in \mathbb{Z} \setminus \{0, \pm 1\}$ , it is true that*

$$(3.37) \quad \langle (L_{c_*} + k^2)\hat{u}_k, \hat{u}_k \rangle_{L^2(\mathbb{R})} \geq A \|\hat{u}_k\|_{H^1(\mathbb{R})}^2,$$

whereas for every  $\hat{u}_0, \hat{u}_{\pm 1} \in H^1(\mathbb{R}) \cap H_\mu^1(\mathbb{R})$ , it is true that

$$(3.38) \quad \langle L_{c_*}\hat{u}_0, \hat{u}_0 \rangle_{L^2(\mathbb{R})} \geq A \|\hat{u}_0\|_{H^1(\mathbb{R})}^2 \quad \text{if } \langle u_{c_*}, \hat{u}_0 \rangle_{L^2(\mathbb{R})} = \langle \partial_\xi^{-1} \partial_c u_c|_{c=c_*}, \hat{u}_0 \rangle_{L^2(\mathbb{R})} = 0$$

and

$$(3.39) \quad \langle (L_{c_*} + 1)\hat{u}_{\pm 1}, \hat{u}_{\pm 1} \rangle_{L^2(\mathbb{R})} \geq A \|\hat{u}_{\pm 1}\|_{H^1(\mathbb{R})}^2 \quad \text{if } \langle \eta_*, \hat{u}_{\pm 1} \rangle_{L^2(\mathbb{R})} = 0.$$

**Proof.** The spectral information on the Schrödinger operator  $L_c : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  with the two lowest eigenvalues (2.4) and (2.5) is sufficient to conclude that  $L_c + k^2 : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is strictly positive for every  $k \in \mathbb{Z} \setminus \{0, \pm 1\}$ . The bound (3.37) follows by the spectral theorem and Gårding's inequality.

Since  $L_{c_*} + 1 : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  is non-negative with a one-dimensional kernel spanned by  $\psi_*$  and  $\langle \eta_*, \psi_* \rangle_{L^2(\mathbb{R})} \neq 0$  by (2.11), this operator is strictly positive under the constraint in (3.39). The constraint in (3.39) is well-defined if  $\hat{u}_{\pm 1} \in H_\mu^1(\mathbb{R})$ . The bound (3.39) follows by the spectral theorem and Gårding's inequality.

Since  $L_{c_*} : H^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$  has one negative and one simple eigenvalues, whereas  $P'(c_*) > 0$ , this operator is non-negative under the first constraint in (3.38) with a one-dimensional kernel spanned by  $\partial_\xi u_{c_*}$  [16]. Since

$$\langle u_{c_*}, \partial_\xi u_{c_*} \rangle_{L^2(\mathbb{R})} = 0, \quad \langle \partial_\xi^{-1} \partial_c u_c|_{c=c_*}, \partial_\xi u_{c_*} \rangle_{L^2(\mathbb{R})} \neq 0,$$

this operator is strictly positive under the two constraints in (3.38). The second constraint in (3.38) is well-defined if  $\hat{u}_0 \in H_\mu^1(\mathbb{R})$ . The bound (3.38) follows by the spectral theorem and Gårding's inequality.  $\blacksquare$

In order to justify the error bound (2.42), we construct the following energy function

$$(3.40) \quad \begin{aligned} F(c) &:= E(u) - E(u_{c_*}) + 4c [P(u) - P(u_{c_*})] \\ &= E(u_0) - E(u_{c_*}) + 4c [P(u_0) - P(u_{c_*})], \end{aligned}$$

where the second equality holds thanks to the conservation of energy  $E$  and momentum  $P$  in time  $t$ . Since  $c(t)$  depends on  $t$ ,  $F(c(t))$  depends on  $t$  but only linearly in  $c(t)$ . By Lemmas 3.1 and 3.2, we rewrite the decompositions (3.2), (3.7), and (3.9) in the form

$$(3.41) \quad u(x, y, t) = u_{c_* + \delta(t)}(\xi) + (b(t)e^{iy} + \bar{b}(t)e^{-iy}) \psi_*(\xi) + v(\xi, y, t).$$

Substituting the decomposition (3.41) into (3.40) yields the following

$$(3.42) \quad F(c) = D(c) + \frac{1}{2} \langle (L_c - \partial_y^2)v, v \rangle_{L^2} + N_c((be^{iy} + \bar{b}e^{-iy})\psi_* + v),$$

where the expansion (3.34) has been used,  $N_c$  satisfies (3.35), and

$$(3.43) \quad \begin{aligned} D(c) &:= E(u_c) - E(u_{c_*}) + 4c [P(u_c) - P(u_{c_*})] \\ &= \frac{1}{2} D''(c_*)(c - c_*)^2 + \tilde{D}(c). \end{aligned}$$

The latter expansion is obtained from  $D(c_*) = D'(c_*) = 0$  and  $D''(c_*) = 4P'(c_*) > 0$  thanks to the variational characterization of the line soliton (2.1) with the action functional (2.2). Thanks to the smoothness of  $D$  in  $c$ , we have  $\tilde{D}(c) = \mathcal{O}((c - c_*)^3)$  as  $c \rightarrow c_*$ .

The following result transfers the initial bound (2.41) to the energy function (3.40).

**Lemma 3.4.** *There exists an  $\varepsilon$ -independent positive constant  $A$  such that*

$$(3.44) \quad |F(c_*)| \leq A\varepsilon^4, \quad |P(u_0) - P(u_{c_*})| \leq A\varepsilon^2.$$

**Proof.** The second bound in (3.44) follows from the initial bound (2.41) thanks to the bound  $|c(0) - c_*| \leq \varepsilon^2$  and the triangle inequality. The first bound in (3.44) follows from the expansion (3.42) with  $D(c_*) = 0$ , the cubic term vanishing

$$(3.45) \quad N_c((be^{iy} + \bar{b}e^{-iy})\psi_*) = 0,$$

the definition  $b(0) = \varepsilon$ , and the triangle inequality. ■

Under the two constraints in (3.3) and the two constraints in (3.8), it follows from Lemma 3.3 that there exists an  $\varepsilon$ -independent constant  $A$  such that

$$(3.46) \quad \langle (L_c - \partial_y^2)v, v \rangle_{L^2} \geq A\|v\|_{H^1(\mathbb{R} \times \mathbb{T})}^2.$$

It follows from the normal form (2.43) that  $|b(t)| \leq C_0\varepsilon$  for every  $t \in \mathbb{R}_+$  as long as  $b(0) = \varepsilon$ . Since  $F(c)$  depends linearly on  $c$  and thanks to (3.46), we obtain the following lower bound from (3.40) and (3.42):

$$\begin{aligned} F(c_*) &\geq 2P'(c_*)(c - c_*)^2 - 4(c - c_*)[P(u_0) - P(u_{c_*})] + \tilde{D}(c) \\ &\quad + \frac{1}{2}A\|v\|_{H^1(\mathbb{R} \times \mathbb{T})}^2 + N_c((be^{iy} + \bar{b}e^{-iy})\psi_* + v). \end{aligned}$$

Thanks to the bounds (3.44) in Lemma 3.4, the smallness of  $N_c$  in (3.35), and the cubic term vanishing in (3.45), we obtain the bound (2.42) for  $|c(t) - c_*|$  and  $\|v\|_{H^1(\mathbb{R} \times \mathbb{T})}$ .

Theorem 1 is proven.

## REFERENCES

- [1] D.C. Bettinson and G. Rowlands, “Transverse stability of plane solitons using the variational method”, *J. Plasma Physics* **59** (1998), 543–554.
- [2] A. Comech, S. Cuccagna, and D. Pelinovsky, “Nonlinear instability of a critical traveling wave in the generalized Korteweg–de Vries equation”, *SIAM J. Math. Anal.* **39** (2007), 1–33.
- [3] R. Cote, C. Munoz, D. Pilod, and G. Simpson, “Asymptotic stability of high-dimensional Zakharov–Kuznetsov solitons”, *Arch. Ration. Mech. Anal.* **220** (2016), 639–710.
- [4] B. Deconinck, D. Pelinovsky, and J.D. Carter, “Transverse instabilities of deep-water solitary waves”, *Proc. Royal Soc. Lond. A* **462** (2006), 2039–2061.
- [5] K.A. Gorshkov and D.E. Pelinovsky, “Asymptotic theory of plane soliton self-focusing in two-dimensional wave media”, *Physica D* **85** (1995), 468–484.
- [6] B.B. Kadomtsev and V.I. Petviashvili, “On the stability of solitary waves in weakly dispersive media”, *Sov. Phys. Dokl.* **15** (1970), 539–541.
- [7] T. Kapitula and A. Stefanov, “A Hamiltonian–Krein (instability) index theory for KdV-like eigenvalue problems”, *Studies in Applied Mathematics* **132** (2014), 183–211.
- [8] T. Kato, *Perturbation theory for linear operators* (Springer–Verlag, Berlin, Heidelberg, 1995).
- [9] Yu.S. Kivshar and D.E. Pelinovsky, “Self-focusing and transverse instabilities of solitary waves”, *Phys. Rep.* **331** (2000), 117–195.
- [10] L.D. Landau and E.M. Lifshitz *Quantum Mechanics* (Volume 3 of *A Course of Theoretical Physics*) (Pergamon Press, New York, 1965).

- [11] D. Lannes, F. Linares, and J.C. Saut, “The Cauchy problem for the Euler–Poisson system and derivation of the Zakharov–Kuznetsov equation”, in *Studies in Phase Space Analysis with Applications to PDEs*, Series Progress in Nonlinear Differential Equations and Applications **84**, M. Cicognani, F. Colombini, D. Del Santo Eds., Birkhäuser, (2013), 183–215.
- [12] F. Linares, A. Pastor, and J.C. Saut, “Well-posedness for the ZK equation in a cylinder and on the background of a KdV soliton”, *Comm. PDEs* **35** (2010), 1674–1689.
- [13] J.L. Marzuola, S. Raynor, and G. Simpson, “Dynamics near the minimal mass soliton for the Korteweg–de Vries equation”, *Dynamical Systems* **29** (2014), 285–299.
- [14] L. Molinet and D. Pilod, “Bilinear Strichartz estimates for the Zakharov–Kuznetsov equation and applications”, *Ann. I.H.Poincaré - AN* **32** (2015), 347–371.
- [15] Y. Murakami and M. Tajiri, “Resonant interaction between  $y$ -periodic soliton and line soliton: solutions to the Kadomtsev–Petviashvili equation with positive dispersion”, *J. Phys. Soc. Japan* **61** (1992), 791–805.
- [16] R.L. Pego and M.I. Weinstein, “Eigenvalues, and instabilities of solitary waves”, *Philos. Trans. Roy. Soc. London A* **340** (1992), 47–94.
- [17] R.L. Pego and M.I. Weinstein, “Asymptotic stability of solitary waves”, *Commun. Math. Phys.* **164** (1994), 305–349.
- [18] D.E. Pelinovsky, “Spectral stability of nonlinear waves in KdV-type evolution equations”, in *Spectral analysis, stability, and bifurcation in modern nonlinear physical systems*, Eds. O. Kirillov and D. Pelinovsky, (John Wiley & Sons, ISTE, 2014), 395–418.
- [19] D.E. Pelinovsky and T.Phan, “Normal form for the symmetry-breaking bifurcation in the nonlinear Schrödinger equation”, *J. Diff. Eqs.* **253** (2012), 2796–2824.
- [20] D.E. Pelinovsky and Yu.A. Stepanyants, “Self-focusing instability of plane solitons and chains of two-dimensional solitons in positive-dispersion media”, *JETF* **77** (1993), 602–608.
- [21] D.E. Pelinovsky, Yu.A. Stepanyants, and Yu.S. Kivshar, “Self-focusing of plane dark solitons in nonlinear defocusing media”, *Phys. Rev. E* **51** (1995), 5016–5026.
- [22] D. E. Pelinovsky, E. A. Ruvinskaya, O. A. Kurkina, B. Deconinck, “Short-wave transverse instabilities of line solitons of the 2-D hyperbolic nonlinear Schrödinger equation”, *Theor. Math. Phys.* **179** (2014), 452–461.
- [23] F. Rousset and N. Tzvetkov, “Transverse nonlinear instability for two-dimensional dispersive models”, *Ann. IHP, Analyse Non Lineaire* **26** (2009), 477–496.
- [24] F. Rousset and N. Tzvetkov, “A simple criterion of transverse linear instability for solitary waves”, *Math. Res. Lett.* **17** (2010), 157–167.
- [25] F. Rousset and N. Tzvetkov, “Stability and instability of the KDV solitary wave under the KP-I flow”, *Commun. Math. Phys.* **313** (2012), 155–173.
- [26] Y. Yamazaki, “Stability of line standing waves near the bifurcation point for nonlinear Schrödinger equations”, *Kodai Math. J.* **38** (2015), 65–96.
- [27] Y. Yamazaki, “Stability for line solitary waves of Zakharov-Kuznetsov equation”, *J. Diff. Eqs.* **262** (2017), 4336–4389.
- [28] A.A. Zaitsev, “Formation of stationary nonlinear waves by superposition of solitons”, *Sov. Phys. Dokl.* **28** (1983), 720–722.
- [29] V.E. Zakharov, “Instability and nonlinear oscillations of solitons”, *JETP Lett.* **22** (1975), 172–173.
- [30] V.E. Zakharov and E.A. Kuznetsov, “On three-dimensional solitons”, *Sov. Phys. JETP* **39** (1974), 285–288.
- [31] V.E. Zakharov and A.M. Rubenchik, “Instability of waveguides and solitons in nonlinear media”, *Sov. Phys. JETP* **38** (1974), 494-500 (1974).

DEPARTMENT OF MATHEMATICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO L8S 4K1, CANADA

DEPARTMENT OF APPLIED MATHEMATICS, NIZHNY NOVGOROD STATE TECHNICAL UNIVERSITY,  
NIZHNY NOVGOROD, RUSSIA

*E-mail address:* dmpeli@math.mcmaster.ca